

/ Please write clearly ir	ո block capitals.	
Centre number	Candidate number	
Surname		
Forename(s)		
Candidate signature	I declare this is my own work.	ノ

A-level **BIOLOGY**

Paper 1

Thursday 10 June 2021

Afternoon

Time allowed: 2 hours

Materials

For this paper you must have:

- a ruler with millimetre measurements
- a scientific calculator.

Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer all questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Show all your working.
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The marks for the questions are shown in brackets.
- The maximum mark for this paper is 91.

For Exam	iner's Use
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
TOTAL	



	Answer all questions in the spaces provided.	
0 1.1	Describe the induced-fit model of enzyme action and how an enzyme acts as a catalyst. [3 marks]
		_
0 1.2	Scientists investigated the action of the enzyme ATP synthase. They made reaction mixtures each containing:	
	 ATP synthase buffer (to control pH) substrates. 	
	One of the substrates required in these reaction mixtures is inorganic phosphate (Pi).	
	Tick (✓) one box to show which other substrate the scientists must add to the reaction mixtures to produce ATP.	
	[1 mark	J
	Adenine	
	Adenosine diphosphate	
	Glucose	
	Ribose	



0 1 . 3 The scientists investigated the effect of concentration of inorganic phosphate (Pi) on ATP synthase activity. After 2 minutes, they stopped each reaction and then measured the concentration of Figure 1 shows the scientists' results. Figure 1 15 10 Concentration of ATP produced / arbitrary units 5 20 10 30 40 50 Concentration of inorganic phosphate (Pi) / mmol dm⁻³ Suggest and explain a procedure the scientists could have used to stop each reaction. [2 marks] 1 Explain the change in ATP concentration with increasing inorganic phosphate concentration. [2 marks]

8



Do not write outside the box


0 2 . 1	Explain the advantage for larger animals of having a specialised system tha	t
	Explain the advantage for larger animals of having a specialised system tha facilitates oxygen uptake.	-
		[2 marks]



Do not write

Figure 2 shows two models of oxygen uptake found in animals.


Figure 2



Oxygen uptake through system developed to the outside of the body, eg fish gills



В



Oxygen uptake through system developed to the inside of the body, eg human lungs

0 2 . 2	Suggest how the environmental conditions have resulted in adaptations of systems
	using Model A rather than Model B.
	[2 mar

[2 marks]		3
	_	

Question 2 continues on the next page






0 2 . 3

Figure 3 shows changes in concentration of oxygen in two gas exchange systems.

Figure 3



Key

- I Air/water entering the gas exchange system
- E Air/water leaving the gas exchange system

Use Figure 3 to justify this conclusion.

- A Arterial blood entering the gas exchange system
- **V** Venous blood leaving the gas exchange system

A student studied **Figure 3** and concluded that the fish gas exchange system is more efficient than the human gas exchange system.

	F
0 2 . 4	Explain how the counter-current principle allows efficient oxygen uptake in the
0 2 . 4	fish gas exchange system.
	[2 marks]



[2 marks]

0 2 . 5

Table 1 shows features of two mammals.

Bats are flying mammals; shrews are ground-living mammals.

Table 1

Mammal Mean body mass / kg		Mean lung volume / cm ³	
Bat	0.096	12.48	
Shrew	0.024	0.72	

Calculate how many times the lung volume per unit of body mass of the bat is greater than that of the shrew.


Give your answer to an appropriate number of significant figures.

Give one suggestion to explain this difference.

[3 marks]

	Answer	
Explanation		
		-







0 3. 1 Describe how one amino acid is added to a polypeptide that is being for ribosome during translation.	ormed at a
noosome during translation.	[3 marks]
-	

Question 3 continues on the next page





Do not write outside the

Table 2 shows:

- mRNA codons and the amino acid coded for by each codon
- the type of bond formed by the R group of some of the amino acids.

Table 2

First	Second base			Third	
base	U	С	Α	G	base
	Phe		Tyr	Cys	U
U		Ser	. , ,		С
	Leu	001	Stop	Stop	Α
	Lou		Отор	Trp	G
			His		U
С	Leu	Pro		Arg	С
			Gln		Α
					G
	lle	Thr	Asn	Ser	U
Α					С
A				Lys	Arg
	Met		Lys	Aig	G
G		Ala	Asp Glu	Gly	U
	Val Ala				С
					Α
		Glu		G	

Key to the type of bond formed by the R group of each amino acid

	Hydrogen bonds Interpretation in the Hydrogen bonds Interpretation District bridges
0 3.2	Crystallin is a structural protein found in the human eye. An inherited disease that leads to blindness is caused by changes in properties of crystallin. The replacement of the amino acid Arg with the amino acid Gly causes these changes. Use information in Table 2 to suggest why this amino acid replacement changes the properties of crystallin. [2 marks]



7

0 3 . 3

The amino acid replacement of Arg with Gly is caused by a single base substitution mutation in the DNA. The non-mutant DNA triplet is TCC.

Complete Table 3.

Give:

- the mRNA codon complementary to the non-mutant DNA triplet
- the mutated mRNA codon that could cause the change from Arg to Gly in the crystallin protein
- the DNA triplet complementary to this mutated mRNA codon.

[2 marks]

Table 3

mRNA codon for the non-mutant triplet	
Mutated mRNA codon	
Mutated DNA triplet	

Turn over for the next question



0 4

A student dissected an organ from a mammal to observe blood vessels.

He dissected a slice of the organ and identified two blood vessels.

Figure 4 shows a photograph of his dissection.

Figure 4





Figure 5 shows a drawing of the blood vessels from his dissection.

Figure 5





	找名校导师 ,
	_
	用小草线上辅导
	肿
	#
	<u>+</u>
	#
	411
	$\overline{}$
_	微信小程序同名
	氚
	\sim
	숲
	냺
	7/
_	쁘
	ПVI

0 4 . 1	Suggest two ways the student could improve the quality of his scientific drawin blood vessels in this dissection.	g of the
	.1	
	2	
0 4.2	Identify the type of blood vessel labelled as X and the type of blood vessel labe Y in Figure 4 .	lled as
	Describe one feature that allowed you to identify the blood vessels.	marks]
	Blood vessel X	
	Blood vessel Y	
	Feature	
0 4.3	Describe two precautions the student should take when clearing away after the dissection.	e marks]
	1	
	2	





Do not write outside the box

0 5.1	Describe how a sample of chloroplasts could be isolated from leaves.	[4 marks]



- 0 5. 2 Scientists grew two groups of plants:
 - control plants with all the inorganic ions needed
 - iron-deficient plants with all the inorganic ions needed **but** without iron ions.

After 1 week, the scientists measured the mass of protein and the mass of chlorophyll in the chloroplasts isolated from samples of leaves of these two groups of plants.

Table 4 shows the scientists' results.

Table 4

Mass of protein / percentage of control	Mass of chlorophyll / percentage of control
40	10

Some proteins found inside the chloroplast are synthesised inside the chloroplast.

Give **one** feature of the chloroplast that allows protein to be synthesised inside the chloroplast **and** describe **one** difference between this feature in the chloroplast and similar features in the rest of the cell.

[2 marks]

Feature
Structural difference

0 5 . 3 The ratio of protein to chlorophyll in control plants is 9:1

Use the information in **Table 4** to calculate the ratio of protein to chlorophyll in iron-deficient plants.

[1 mark]


Question 5 continues on the next page



0 5.4	The scientists also observed the chloroplasts from the samples of leaves using an electron microscope.	
	Figure 6 shows a chloroplast from a control plant (image A) and a chloroplast from an iron-deficient plant (image B).	
	Figure 6	
	This source has been removed due to third-party copyright restrictions.	
	Use Figure 6 to suggest why iron-deficient plants have a reduced growth rate. [3 marks]	



Do not write outside the








0 6 . 1

Figure 7 shows the mean distance between centromeres and the poles (ends) of the spindle during mitosis.

Figure 7



Calculate the rate of movement of the centromeres during phase E.

Give your answer in µm minute⁻¹ and to 3 decimal places.

[2 marks]

um	minute-1
мии	minato



找名校导师,用小草线上辅导(微信小程序同名)

0 6.2	Name the three phases of mitosis shown by C , D and E on Figure 7 .
	Describe the role of the spindle fibres and the behaviour of the chromosomes during each of these phases.
	[5 marks]
	c
	D
	E

Turn over for the next question




0 7 . 1

ADCs are molecules made of a monoclonal antibody linked to a cancer drug.

Figure 8 shows how an ADC enters and kills a tumour cell.

The process of entering the cell and the breakdown of the antibody to release the drug is very similar to phagocytosis.

Figure 8



Use your knowledge of phagocytosis to describe how an ADC enters and kills the tumour cell.



[3 marks]

0 7.2	Some of the antigens found on the surface of tumour cells are also found on the surface of healthy human cells.
	Use this information to explain why treatment with an ADC often causes side effects. [2 marks]

Question 7 continues on the next page



Scientists investigated whether one type of ADC could be used to treat human breast cancer.

This ADC is a monoclonal antibody combined with a drug to inhibit mitosis. The monoclonal antibody binds to a protein found on human breast cancer cells.

The scientists placed small pieces of human breast cancer tissue under the skin of mice.

The scientists then randomly divided the mice into three groups. They treated the groups as follows on day 0.

Group G - control

Group H – injected with monoclonal antibody only

Group J – injected with ADC (monoclonal antibody + drug).

Every few days, the scientists measured the volume of the tumours formed from the human breast cancer tissue.

Figure 9 shows the scientists' results.

Figure 9

This source has been removed due to third-party copyright restrictions.



0 7.3	Mice in Group H were injected with 2 mg kg ⁻¹ of monoclonal antibody. The monoclonal antibody was in a solution of concentration 500 mg dm ⁻³	(
	Calculate the volume of antibody solution that the scientists would have injected into a 23 g mouse. Give your answer in dm³ and in standard form. [2 marks]	
	dm ³	
0 7.4	Suggest one reason why there are no data for Group G and Group H after day 8 [1 mark]	
0 7.5	Suggest and explain two further investigations that should be done before this ADC is tested on human breast cancer patients.	
	[2 marks]	
	2	
		_

Turn over for the next question



Do not write outside the box

0 8 . 1	Describe how a triglyceride molecule is formed.	[3 marks]



0 8 . 2

Table 5 shows some properties of four fatty acids.

Table 5

Fatty acid	Number of carbon atoms in the R group	Number of double bonds in the R group
Caprylic acid	8	0
Palmitoleic acid	16	1
Stearic acid	18	0
Linoleic acid	18	2

Figure 10 shows diagrams of these fatty acids.

Figure 10

Put a tick (\checkmark) in **one** box that contains correct information about one of these fatty acids.

[1 mark]

Caprylic acid is an unsaturated fatty acid represented by diagram L .	
Linoleic acid is a saturated fatty acid represented by diagram N .	
Palmitoleic acid is an unsaturated fatty acid represented by diagram K .	
Stearic acid is a saturated fatty acid represented by diagram M .	



The percentage of saturated fatty acids compared with unsaturated fatty acids found in lipid stores in seeds differs in different populations.

Scientists investigated two populations of the plant, *Helianthus annuus*.

The scientists grew young plants from seeds collected from each population. They placed the seeds on wet tissue paper so that the root growth was visible.

They grew seeds from each population at two temperatures:

- warm temperature of 24 °C
- cool temperature of 10 °C

After 10 days, the scientists measured the length of each root.

Table 6 shows some of the properties of the two populations and the scientists' results.

Table 6

Population	Temperature in natural environment	In the seed – Mean percentage of fatty acids that are saturated	Mean length of root after 10 days at 24°C / mm (± 2 x standard deviation)	Mean length of root after 10 days at 10 °C / mm (± 2 x standard deviation)
1	Warm	10.9	8.2 (±1.0)	3.1 (±0.3)
2	Cool	6.1	5.5 (±0.9)	4.3 (±0.2)


The mean ±2 × standard deviation includes 95% of the data.

0 8 . 3	The scientists used a data logger to measure the length of the root rather than a ruler.
	Suggest one reason why they used a data logger and explain why this was important in this investigation.
	[1 mark]



8 . 4	It is known that:
	during respiration saturated fatty acids yield more energy than unsaturated fatty acids
	 saturated fatty acids have higher melting points than unsaturated fatty acids lipases in seeds act more rapidly on liquid substrates.
	Use this information and Table 6 to show how each population is better adapted for its natural environment when compared with the other population.
	[4 marks]
8 . 5	Although these two populations are completely separate and show genetic variation, they are both called <i>Helianthus annuus</i> .
	Explain why they are both given this name. [1 mark]







Omple

Complete **Table 7** with ticks (\checkmark) to show which elements are found in the following biological molecules.

[2 marks]

Table 7

Biological molecules	Element				
	Carbon	Nitrogen	Oxygen	Phosphorus	
Galactose					
Phospholipid					
RNA					
Sucrose					

Question 9 continues on the next page

2 9


After Watson and Crick proposed the model of DNA structure, scientists investigated the possible mechanisms for DNA replication.

Two scientists grew a bacterial population, providing them with a nitrogen source containing only the heavy isotope of nitrogen, ¹⁵ N. As soon as all the DNA in this population contained ¹⁵ N, the scientists changed the nitrogen source to one containing only the lighter isotope of nitrogen, ¹⁴ N. They changed the nitrogen source at 0 hours.

During the investigation, the scientists measured the size of the population of bacterial cells.

Figure 11 shows the scientists' results.







0 9 . 2

The generation time for a population of bacteria is the time taken for all the bacteria to divide once by binary fission.

Use **Figure 11** and the following equation to calculate the generation time for this population of bacteria. Give your answer in hours.

[2 marks]

Number of generations =
$$\frac{\log_{10} \left(\frac{\text{size of population at time } + 4 \text{ hours}}{\text{size of population at time } - 4 \text{ hours}} \right)}{\log_{10} 2}$$

Generation time hours

Question 9 continues on the next page



At intervals during this investigation, the scientists removed samples of the bacterial population, isolated the DNA and measured the density of the DNA.

Figure 12

DNA made using ¹⁵N has a higher density than DNA made using ¹⁴N.

Figure 12 shows the scientists' results.

Relative concentration of DNA


Density

Generation 0

Generation 0

Generation 1

Generation 1



Relative concentration of DNA

Density

Generation 3





7


There are **three** possible models of DNA replication.

These models are shown in Figure 13.

Figure 13







0 9 . 3 Which of these models, P, Q or R, is supported by the results shown in Figure 12?

Give the letter and name of the model supported and explain why the results do **not** support the other models.

[3 marks]

Model		
Model		

Name

Explanation for first unsupported model

Explanation for second

unsupported model



Do not write outside the box

1 0.1	Describe the structure of DNA. [5 mail	rks]



1 0.2	Name and describe five ways substances can move across the cell-surface membrane into a cell.	
		[5 marks]
	Question 10 continues on the next page	






Figure 14 shows transmission electron micrographs of two cells, one animal cell and one prokaryotic cell.


Figure 14

Cell A



Magnification ×30 000

Cell B




Magnification ×60 000



0 . 3	Contrast the structure of the two cells visible in the electron micrographs shown in Figure 14 .	outside box
	[5 marks]	
		-
		-
		-
		-
		-
		-
		-
		-
		-
		-
		-
		-
		-
		-
		-
		-
		-
		_
		15
	END OF QUESTIONS	







Question number	Additional page, if required. Write the question numbers in the left-hand margin.



Question number	Additional page, if required. Write the question numbers in the left-hand margin.
	Copyright information
	For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from www.aqa.org.uk.
	Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.
	Copyright © 2021 AQA and its licensors. All rights reserved.



