

Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

Forename(s)

Candidate signature

I declare this is my own work.

A-level MATHEMATICS

Paper 2

Time allowed: 2 hours

Materials

- You must have the AQA Formulae for A-level Mathematics booklet.
- You should have a graphical or scientific calculator that meets the requirements of the specification.

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer each question in the space provided for that question. If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do **not** write outside the box around each page or on blank pages.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 100.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	
16	
17	
18	
19	
TOTAL	



Section A

Answer **all** questions in the spaces provided.

- 1** A circle has centre $(4, -5)$ and radius 6

Find the equation of the circle.

Tick (✓) **one** box.**[1 mark]**

$$(x - 4)^2 + (y + 5)^2 = 6$$

☐

$$(x + 4)^2 + (y - 5)^2 = 6$$

☐

$$(x - 4)^2 + (y + 5)^2 = 36$$

☐

$$(x + 4)^2 + (y - 5)^2 = 36$$

☐

- 2** State the value of

$$\lim_{h \rightarrow 0} \frac{\sin(\pi + h) - \sin \pi}{h}$$

Circle your answer.

[1 mark] $\cos h$

-1

0

1







3 The function f is concave and is represented by one of the graphs below.

Identify the graph which represents f .

Tick (✓) **one** box.

[1 mark]


☐

☐

☐

☐

Turn over ►



Find Personal Tutor from www.wisesprout.co.uk

找名校导师，用小草线上辅导（微信小程序同名）

A triangle with vertices labeled A, B, and C. Vertex A is at the top, B is at the bottom left, and C is at the bottom right. The side AC is labeled 6.1 cm. The side BC is labeled 8.7 cm. The interior angle at vertex B is labeled 38°.

AB is the shortest side. The lengths of AC and BC are 6.1 cm and 8.7 cm respectively.

Find the size of the largest angle.

Give your answer to the nearest degree.

[3 marks]

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page. There are no margins, text, or other markings on the paper.

Turn over for the next question

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Find Personal Tutor from www.wisesprout.co.uk

找名校导师，用小草线上辅导（微信小程序同名）

Turn over ►



0 5

5 (c) Hence, or otherwise, find

$$\int \left((2 + 5x)^4 - (2 - 5x)^4 \right) dx$$

[2 marks]

Turn over for the next question

Turn over ►



6 (a) Asif notices that $24^2 = 576$ and $2 + 4 = 6$ gives the last digit of 576

He checks two more examples:

$$27^2 = 729$$

$$2 + 7 = 9$$

Last digit 9

$$29^2 = 841$$

$$2 + 9 = 11$$

Last digit 1

Asif concludes that he can find the last digit of any square number greater than 100 by adding the digits of the number being squared.

Give a counter example to show that Asif's conclusion is **not** correct.

[2 marks]

6 (b) Claire tells Asif that he should look only at the last digit of the number being squared.

$$27^2 = 729$$

$$7^2 = 49$$

Last digit 9

$$24^2 = 576$$

$$4^2 = 16$$

Last digit 6

Using Claire's method determine the last digit of 23456789^2

[1 mark]



- 6 (c)** Given Claire's method is correct, use proof by exhaustion to show that no square number has a last digit of 8


[2 marks]

Turn over for the next question

Turn over ►



The curve $y = 15 - x^2$ and the isosceles triangle OPQ are shown on the diagram below.



The line PQ is parallel to the x -axis.

Show that the area, A , of the triangle OPQ is given by

$$A = 15q - q^3 \quad \text{for } 0 < q < c$$

where c is a constant to be found.

[3 marks]

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Fully justify your answer.

[6 marks]

[illegible]


Turn over for the next question

Turn over ►



8 (a) Sketch the graph of $y = \frac{1}{x^2}$

[2 marks]



- 8 (b)** The graph of $y = \frac{1}{x^2}$ can be transformed onto the graph of $y = \frac{9}{x^2}$ using a stretch in one direction.

Beth thinks the stretch should be in the y -direction.

Paul thinks the stretch should be in the x -direction.

State, giving reasons for your answer, whether Beth is correct, Paul is correct, both are correct or neither is correct.

[3 marks]

Turn over for the next question

Turn over ►



Given that

show that

$$x = Ay^p$$

where A is an integer and p is a rational number.

[4 marks]

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Turn over for the next question

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Find Personal Tutor from www.wisesprout.co.uk

找名校导师，用小草线上辅导（微信小程序同名）

Turn over ►



1 5

10 A gardener has a greenhouse containing 900 tomato plants.

The gardener notices that some of the tomato plants are damaged by insects.

Initially there are 25 damaged tomato plants.

The number of tomato plants damaged by insects is increasing by 32% each day.

10 (a) The total number of plants damaged by insects, x , is modelled by

$$x = A \times B^t$$

where A and B are constants and t is the number of days after the gardener first noticed the damaged plants.

10 (a) (i) Use this model to find the total number of plants damaged by insects 5 days after the gardener noticed the damaged plants.

[3 marks]

10 (a) (ii) Explain why this model is not realistic in the long term.

[2 marks]




Find Personal Tutor from www.wisesprout.co.uk

找名校导师，用小草线上辅导（微信小程序同名）

[3 marks]

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page. There are no margins, text, or other markings on the paper.

Turn over ►



[5 marks]

[illegible]

[2 marks]



Section B

Answer **all** questions in the spaces provided.

11

A moon vehicle has a mass of 212 kg and a length of 3 metres.

On the moon the vehicle has a weight of 345 N

Calculate a value for acceleration due to gravity on the moon.

Circle your answer.

[1 mark]

0.614 m s^{-2}

1.63 m s^{-2}

1.84 m s^{-2}

4.89 m s^{-2}

12

A car is travelling along a straight horizontal road with initial velocity $u \text{ m s}^{-1}$

The car begins to accelerate at a constant rate $a \text{ m s}^{-2}$ for 5 seconds, to reach a final velocity of $4u \text{ m s}^{-1}$

Express a in terms of u .

Circle your answer.

[1 mark]

$a = 0.2u$

$a = 0.4u$

$a = 0.6u$

$a = 0.8u$

Turn over for the next question

Turn over ►

A ball is projected from a point on horizontal ground with an initial velocity of 7 m s^{-1} at an angle θ above the horizontal.

13 (a) Show that

$$h = 2.5 \sin^2 \theta$$

[3 marks]

[illegible]

13 (b) Hence, given that $0^\circ \leq \theta \leq 60^\circ$, find the maximum value of h .

[2 marks]



- 13 (c)** Nisha claims that the larger the size of the ball, the greater the maximum vertical height will be.

State whether Nisha is correct, giving a reason for your answer.

[1 mark]

Turn over for the next question

Turn over ►



- 14** A £2 coin has a diameter of 28 mm and a mass of 12 grams.

A uniform rod AB of length 160 mm and a fixed load of mass m grams are used to check that a £2 coin has the correct mass.

The rod rests with its midpoint on a support.

A £2 coin is placed face down on the rod with part of its curved edge directly above A .

The fixed load is hung by a light inextensible string from a point directly below the other end of the rod at B , as shown in the diagram.



- 14 (a)** Given that the rod is horizontal and rests in equilibrium, find m .

[3 marks]


- 14 (b)** State an assumption you have made about the £2 coin to answer **part (a)**.

[1 mark]



A car is moving in a straight line along a horizontal road.

Find Personal Tutor from www.wisesprout.co.uk



找名校导师，用小草线上辅导（微信小程序同名）

Initially the car has velocity 0 m s^{-1}

Find the next time when the velocity of the car is 0 m s^{-1}

[4 marks]

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page. There are no margins, text, or other markings on the paper.

Turn over ►



Particle P is initially at rest at the point with position vector $(-4\mathbf{i} + 5\mathbf{j})$ metres and moves with constant acceleration $(3\mathbf{i} - 4\mathbf{j}) \text{ m s}^{-2}$

P and Q are moving along parallel paths.

[4 marks]

This image shows a single sheet of white paper with horizontal blue or grey ruling lines. The lines are evenly spaced and run across the width of the page. There are approximately 20 lines visible. The paper has a slight shadow on the right side, suggesting it's resting on a surface.

[1 mark]



[3 marks]

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page. There are no margins, text, or other markings on the paper.

Find Personal Tutor from www.wisesprout.co.uk

找名校导师，用小草线上辅导（微信小程序同名）

Turn over ►



A particle is moving such that its position vector, $\mathbf{r}$ metres, at time t seconds, is given by

Show that the **magnitude** of the acceleration of the particle, $a \text{ m s}^{-2}$, is given by

Fully justify your answer.

[illegible]

Turn over for the next question

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Find Personal Tutor from www.wisesprout.co.uk

找名校导师，用小草线上辅导（微信小程序同名）

Turn over ►



An object, O, of mass m kilograms is hanging from a ceiling by two light, inelastic strings of different lengths.

The longer string, of length 1.2 metres, is fixed to the ceiling at B .

Show that the tension in the shorter string is over 30% more than the tension in the longer string.

[4 marks]

[illegible]

Find the value of m .

This image shows a blank sheet of white paper with horizontal blue ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.


Turn over ►



In this question use $g = 9.8 \text{ m s}^{-2}$

A crate of mass 20 kg is at rest at the point A on the ramp.

Once in motion, the rope remains taut and parallel to the line of greatest slope of the ramp.



The tension in the rope is 230 N

Find the coefficient of friction between the crate and the ramp.

[7 marks]

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

19 (b) (i) The crate takes 3.8 seconds to reach the top of the ramp.

Find the distance OA .

[3 marks]

19 (b) (ii) Other than air resistance, state **one** assumption you have made about the crate in answering part **(b)(i)**.

[1 mark]

END OF QUESTIONS



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Find Personal Tutor from www.wisesprout.co.uk

找名校导师，用小草线上辅导（微信小程序同名）



[illegible]

[illegible]

[illegible]

