

Please write clearly in	า block capitals.
Centre number	Candidate number
Surname	
Forename(s)	
Candidate signature	I declare this is my own work.

GCSE CHEMISTRY



Higher Tier Paper 2

Tuesday 13 June 2023 Morning Time allowed: 1 hour 45 minutes

Materials

For this paper you must have:

- a ruler
- · a scientific calculator
- the periodic table (enclosed).

Instructions

- Use black ink or black ball-point pen.
- · Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer all questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do all rough work in this book. Cross through any work you do not want to be marked.
- In all calculations, show clearly how you work out your answer.

Information

- The maximum mark for this paper is 100.
- The marks for questions are shown in brackets.
- You are expected to use a calculator where appropriate.
- You are reminded of the need for good English and clear presentation in your answers.

For Exam	iner's Use
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
TOTAL	ľ




0 1

A student investigated the rate of the reaction between zinc and sulfuric acid.

Hydrogen gas is produced during this reaction.

Figure 1 shows the apparatus.

Figure 1



This is the method used.

- 1. Add 50 cm³ of sulfuric acid to a conical flask.
- 2. Add 2.0 g of zinc to the conical flask.
- 3. Quickly put a stopper in the conical flask and start a timer.
- 4. Measure the time taken to collect 20 cm³ of gas.
- 5. Repeat steps 1 to 4 three more times.

0 1 . 1	Suggest why the stopper must be put in the conical flask as quickly as possible in step 3 .
	[1 mark]



0 1.2 The student calculated the rate of the reaction for each trial.

Table 1 shows the results of the calculations.

Table 1

	Trial 1	Trial 2	Trial 3	Trial 4
Rate of reaction in cm ³ /s	0.78	0.81	0.68	0.81

Determine the mean time taken to collect 20 cm³ of gas.			
Do not include an	y anomalous results.		
Use the equation:			
	mean rate of reaction =	volume of gas collected mean time taken	
		mean time taken	[5 marks]

Mean time taken = _____

Question 1 continues on the next page




Turn over ►

	找名校导师,
	用小草线上辅导
	$\overline{}$
-	微信小程序同名

10

0 1.3	The student changed the investigation so that the mean time taken to collect 20 cm ³ of gas was greater.	
	Which two changes would increase the mean time taken to collect 20 cm ³ of gas? [2 marks]	
	Tick (✓) two boxes.	
	Use a catalyst	
	Use a larger conical flask	
	Use a lower temperature	
	Use smaller pieces of zinc	
	Use sulfuric acid of a lower concentration	
0 1.4	Hydrogen gas is produced during this reaction. Describe the test for hydrogen gas.	
	Give the result of the test. [2 marks]	
	Test	
	Result	









0 2

This question is about alcohols and carboxylic acids.

Alcohols are used as fuels.

A student burned 1.00 g of six alcohols and determined the energy released from each.

Table 2 shows the results.

Table 2

Alcohol	Formula of one molecule of the alcohol	Energy released in kJ/g
Ethanol	C ₂ H ₅ OH	29.6
Propanol	C₃H ₇ OH	33.6
Butanol	C ₄ H ₉ OH	36.1
Pentanol	C₅H ₁₁ OH	37.7
Hexanol	C ₆ H ₁₃ OH	38.9
Heptanol	C ₇ H ₁₅ OH	39.8

0 2 . 1		Calculate the mass of ethanol that must be burned to release the sa	ime amount of
		energy as burning 1.00 g of heptanol.	[2 marks]
		Mass =	9


0 2.2 The energy released in kJ/g varies with the number of carbon atoms in one molecule of each alcohol.

Plot the data from **Table 2** on **Figure 2**.

[2 marks]







0 2.3 Estimate the energy released in kJ when 1.00 g of octanol (C₈H₁₇OH) is burned. Use **Figure 2**.

[1 mark]

Energy released = kJ

Turn over ▶



	Carbon dioxide is produced when alcohols are burned. Carbon dioxide is identified by bubbling the gas through limewater.	
0 2.4	Complete the sentence.	
	Choose the answer from the box.	[1 mark]
	calcium chloride calcium hydroxide calcium nitrate calcium s	sulfate
	Limewater is an aqueous solution of	
0 2.5	Give the result of the test when carbon dioxide is bubbled through limewater.	[1 mark]




	Ethanoic acid can be produced from ethanol.	Do not write outside the box
0 2 . 6	What is reacted with ethanol to produce ethanoic acid? Tick (✓) one box. [1 mark]	
	A halogen	
	An alkali metal	_
	An oxidising agent	חוום רפוסטומ
	Water	
0 2.7	Ethanoic acid contains the functional group –COOH	www.wasaprodi.co.ds
	Complete the displayed structural formula of this functional group. [1 mark]	
		3.4 世 文 4 岩 -
	-c o	137 1 %
	о — H	
		1.7 VE/17/19 11
		,
	Question 2 continues on the next page	



10 Do not write outside the 0 2 . 8 Ethanoic acid reacts with different compounds. Draw one line from each compound to a product of the reaction of the compound with ethanoic acid. [2 marks] Product of the reaction Compound with ethanoic acid Carbon dioxide Ethene Ethanol Ethyl ethanoate Sodium carbonate Hydrogen Poly(ethene)









Do not write outside the box

0 3	This question is about chemical analysis.
	Potassium bromide is used in medicine.
	A scientist tested a sample of medicine to show the presence of potassium ions and of bromide ions.
	The sample is soluble in water.
0 3.1	Plan a method the scientist could use to show that the sample of medicine contains potassium ions and bromide ions.
	The scientist has:
	a Bunsen burner
	a metal wire
	• test tubes
	a dropping pipette
	distilled water
	dilute nitric acid
	silver nitrate solution.
	You should give the results of the tests. [6 marks]



8

找名校导师,
用小草线上辅导
(微信小程序同名)

Do not write The scientist could also use an instrumental method to show the presence of potassium ions in the medicine. 0 3 . Which instrumental method could be used to show the presence of potassium ions in the medicine? [1 mark] 3 Give **one** advantage of using this instrumental method instead of a chemical test. [1 mark]

Turn over for the next question



Do not write outside the box

0 4	This question is about greenhouse gases and climate change.	
	Carbon dioxide and methane are greenhouse gases.	
0 4.1	Which of the following is also a greenhouse gas?	[1 mark]
	Tick (✓) one box.	
	Chlorine	
	Nitrogen	
	Oxygen	
	Water vapour	
	In the past 50 years, there has been an increase in:	
	the world population	
	the concentration of carbon dioxide in the atmosphere	
	the concentration of methane in the atmosphere	
	the mean temperature of the atmosphere at the Earth's surface.	
	Most scientists think this information can be used to explain climate change.	
0 4.2	Explain why the increase in world population may have caused the increase in concentration of carbon dioxide in the atmosphere.	n the
		2 marks]



0 4 . 3	Explain why the increase in world population may have caused the increase in the	Do not write outside the box
	concentration of methane in the atmosphere.	
	[2 marks]	
		7
		5
		<u> </u>
0 4 . 4	Describe two potential effects of the increase in the mean temperature of the	۵
	atmosphere at the Earth's surface. [2 marks]	
	1	*
		T.WIO
		<u>.</u>
	2	wasabiodi.co.dx
		5
		Į Ž
		\$ 4
		,
0 4 . 5	The mean temperature of the atmosphere at the Earth's surface has increased.	<u>ئ</u> د نا
	Most esigntists think that this has been equeed by an increase in the concentration of	л И Н
	Most scientists think that this has been caused by an increase in the concentration of greenhouse gases in the atmosphere.	± 4
		- الله الله
	Give one reason why some scientists do not accept this theory.	j
	[1 mark]	1±/J7
		1 1 1 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
		8
	Turn over for the next question	
	o to to to the modern	



0 5	Copper is extracted from metal ores. Chalcopyrite is a metal ore containing a compound with the formula CuFeS ₂
0 5.1	CuFeS $_2$ reacts with oxygen to produce copper(II) sulfate and iron(II) sulfate. Complete the equation for this reaction. You should balance the equation. [2 marks]
	CuFeS ₂ + \rightarrow CuSO ₄ + FeSO ₄
0 5.2	Calculate the percentage by mass of copper in $CuFeS_2$ Relative atomic masses (A_r): $S = 32$ Fe = 56 Cu = 63.5 [3 marks]
0 5.3	Describe a test to show the presence of copper(II) ions in a solution of copper(II) sulfate. Give the result of the test.
	Test
	Result



0 5.4	Copper can be extracted from low-grade ores by bioleaching.		outside th
	Describe what is meant by bioleaching.	[2 marks]	
			9
			1

Turn over for the next question

Turn over ▶



Do not write

0 6

This question is about chromatography.

A student investigated an orange food colouring using two different types of chromatography paper.

The food colouring:

- · contained a mixture of red and yellow dyes
- was soluble in water.

This is the method used.

- 1. Draw a start line on a piece of type **A** chromatography paper.
- 2. Put a spot of orange food colouring on the line.
- 3. Put the paper into a beaker containing water as a solvent.
- 4. Wait for the water to travel up the paper.
- 5. Measure the distance above the start line moved by the red and yellow dyes and the water.
- 6. Repeat steps 1 to 5 using type **B** chromatography paper.




Do not write outside the

box

Figure 3 shows how the student set up the apparatus.





0	6		1	The student made two mistakes when setting up the apparatus.
---	---	--	---	---

Give two mistakes the student made.

[2 marks]

1			
2			

Question 6 continues on the next page



Another student set up the apparatus correctly.

Table 3 shows the results.

Table 3

	Type A chromatography paper		Type B chromatography pape	
	Red dye	Yellow dye	Red dye	Yellow dye
Distance moved by dye in cm	4.8	6.6	5.4	х
Distance moved by water in cm	12.0	12.0	12.0	12.0
R _f value	0.40	0.55	0.45	0.60

0	6 . 2	Determine value X in Table 3 .		[3 marks]
			X =	cm



Find Personal Tutor from www.wisesprout.co.uk
Tutor fro
rom www.v
wisesprou
it.co.uk

找名校导师,用<u>小草线上辅</u>导(微信小程序同名)

	Changing the type of chromatography paper resulted in different R_{f} values for the red dye.
0 6.3	Explain why the $R_{\mbox{\scriptsize f}}$ values for the red dye are different using the two types of chromatography paper.
	Use Table 3 . [3 marks]
0 6.4	What other change to the investigation could result in a different R_f value for the
	red dye? [1 mark]

Turn over for the next question



Turn over ▶

Do not write outside the box

0 7	Manganese dioxide catalyses the decomposition of hydrogen peroxide solution.
	Oxygen and water are produced.
0 7.1	Explain how a manganese dioxide catalyst increases the rate of decomposition of hydrogen peroxide. [2 marks]
	A student investigated the rate of this reaction.
	This is the method used.
	1. Add 50 cm ³ of 2.0 mol/dm ³ hydrogen peroxide solution to a conical flask.
	2. Add 1.0 g of manganese dioxide to the conical flask.
	3. Place the conical flask on a balance and start a timer.
	4. Record the total mass lost from the conical flask every 20 seconds for 180 seconds.
0 7.2	Explain why the mass of the conical flask and contents decreased.
	[2 marks]




0 7.3

Figure 4 shows the results for 50 cm³ of 2.0 mol/dm³ hydrogen peroxide solution and 1.0 g of manganese dioxide.

A tangent to the line has been drawn at 75 seconds.

Figure 4



Determine the rate of reaction when the time was 75 seconds.

Give your answer to 2 significant figures.

[4 marks]

Rate (2 significant figures) =	g/s
--------------------------------	-----


Turn over ▶



0 7.4

The results for 50 cm³ of 2.0 mol/dm³ hydrogen peroxide solution and 1.0 g of manganese dioxide are shown again on **Figure 5**.

Figure 5



The student repeated the investigation using 50 cm³ of 1.0 mol/dm³ hydrogen peroxide solution and 1.0 g of manganese dioxide.

Sketch the expected results for 1.0 mol/dm³ hydrogen peroxide solution on **Figure 5**. **[2 marks]**




找名校导师,用小草线上辅导(微信小程序同名)

This question is about polymers.

Chloroethene can be used to produce an addition polymer called poly(chloroethene).

The displayed structural formula of chloroethene is




Draw a circle around the functional group on the displayed structural formula that allows chloroethene to produce an addition polymer.

[1 mark]

0 8 . 2 Complete the equation for the production of poly(chloroethene) from chloroethene.

[3 marks]



0 8 . 3 Poly(ethene) can be strengthened with wood particles to make a building material.

The building material consists of a wood particle reinforcement embedded in a poly(ethene) matrix.

What general name is given to materials like this?

[1 mark]



0 8.4	The amino acid beta-alanine has the formula
	H ₂ NCH ₂ CH ₂ COOH
	Beta-alanine polymerises to produce a polypeptide and a small molecule.
	Name the small molecule produced when beta-alanine polymerises. [1 mark]
0 8 . 5	An amino acid can be represented as:
	H_2N — COOH
	The relative formula mass (M_r) of this amino acid is 75
	Calculate the relative formula mass of the section of this amino acid molecule represented by
	Relative atomic masses (A_r): H = 1 C = 12 N = 14 O = 16 [2 marks]

Relative formula mass =



Figure 6 represents part of a naturally occurring polymer molecule produced from glucose.

Figure 6

0 8. 6 Draw a circle around the repeating unit in the polymer in Figure 6.


[1 mark]

0 8 . 7 Suggest the identity of this polymer.

[1 mark]

Figure 7 represents the structure of a naturally occurring polymer.

Figure 7



0 8 . 8 Give the general name for the four different monomers which make up the structure shown in **Figure 7**.

[1 mark]


0 8 . 9 Name the **shape** of the structure shown in **Figure 7**.

[1 mark]

12

Turn over ▶







0 9	This question is about reversible reactions.
	 When 4.68 g of hydrated copper sulfate changes into anhydrous copper sulfate: 2.99 g of anhydrous copper sulfate is produced 1.47 kJ of energy is taken in from the surroundings.
	The equation for the reversible reaction is:
	hydrated copper sulfate ⇌ anhydrous copper sulfate + water
0 9.1	Calculate the maximum mass of water that can be produced from 11.7 g of hydrated copper sulfate. [3 marks]
	Mass = g
0 9.2	15.0 g of anhydrous copper sulfate completely changes into hydrated copper sulfate when water is added.
	Calculate the amount of energy transferred to the surroundings. [2 marks]
	Energy =kJ





	The gases nitrogen dioxide and dinitrogen tetroxide reach dynamic equilibrium in a sealed container.		
	The equation for the reaction is:		
	2 NO ₂ (g)	\rightleftharpoons	$N_2O_4(g)$
	nitrogen dioxide (brown)		dinitrogen tetroxide (colourless)
	The forward reaction is exothermic.		
0 9 . 3	What happens to the position of the equis increased?	uilibrium	
	Tick (✓) one box.		[1 mark]
	Shifts to the left		
	Stays the same		
	Shifts to the right		



	31
0 9.4	A teacher seals a brown-coloured mixture of nitrogen dioxide and dinitrogen tetroxide in a gas syringe.
	Figure 8 shows the sealed gas syringe.
	Figure 8
	Syringe piston Seal Mixture of gases
	The teacher pushes the syringe piston in.
	This increases the pressure in the gas syringe.
	What is the colour of the mixture when a new equilibrium position is reached? [1 mark] Tick (✓) one box.
	The mixture is a darker shade of brown.
	The mixture is the same shade of brown.
	The mixture is a lighter shade of brown.

Question 9 continues on the next page



Hydrogen iodide gas decomposes into hydrogen gas and iodine gas at high temperatures.

The equation for the reaction is:

$$2 HI(g) \Rightarrow H_2(g) + I_2(g)$$

0 9 . 5	Explain the effect of increasing the pressure on the equilibrium position of this reaction.	
		[2 marks]

0 9 . 6	Suggest the effect of adding a catalyst on the equilibrium position of this reaction.
	[1 mark]



		Do not write
	Copper forms coloured compounds.	outside the
	Hydrochloric acid is added to an aqueous solution of copper compound A .	
	The word equation for the reaction is:	
	copper compound A + hydrochloric acid	
0 9 . 7	The reaction mixture is green when both copper compounds are present in a solution	
0 9 . 7	The reaction mixture is green when both copper compounds are present in a solution at equilibrium.	
	How can the equilibrium position be shifted to make the reaction mixture more yellow? [1 mark]	
	Tick (✓) one box.	
	Add more hydrochloric acid	
	Add more water	
	Leave the reaction mixture for 30 minutes	
0 9.8	The concentrations of the substances in this reaction do not change at dynamic equilibrium.	
	Explain why. [2 marks]	
		13





Do not write

1 0

This question is about fertilisers.

Compounds of nitrogen (N), phosphorus (P) and potassium (K) are used as fertilisers to improve agricultural productivity.

Table 4 shows information about three compounds, $\bf A$, $\bf B$ and $\bf C$, that can be used as fertilisers.

Table 4

	Compound A	Compound B	Compound C	
Name	potassium chloride	ammonium nitrate	diammonium hydrogen phosphate	
Formula	KCl	NH ₄ NO ₃	(NH ₄) ₂ HPO ₄	
Percentage (%) of N, P and K by mass	K : 52%	N : 35%	N: 21% P: 23%	
Cost in £/kg	0.24	0.23	0.35	

1 0 . 1	A scientist analysed the percentages of nitrogen, phosphorus and potassium in a soil.
	The percentages of nitrogen and of potassium in the soil were lower than the percentages needed for high agricultural productivity.
	There was sufficient phosphorus in the soil for high agricultural productivity.
	Evaluate the use of the compounds in Table 4 to improve the agricultural productivity of this soil.
	[4 marks]


-		
-		
-		



10.2	How is potassium chloride (compound A) obtained from the Earth?	[1 mark]	Do not writ outside the box
1 0 . 3	Name one other compound that could be used instead of potassium chloride (compound A) to give a similar improvement in agricultural productivity.	[1 mark]	
10.4	Nitric acid is needed to produce ammonium nitrate (compound B). Name a compound needed to produce nitric acid.	[1 mark]	
1 0.5	Phosphate rock contains phosphorus compounds. Plants absorb phosphorus from compounds dissolved in rainwater. Suggest why phosphate rock cannot be used directly as a fertiliser.	[1 mark]	
1 0 . 6	Phosphate rock can be treated with different acids to produce salts useful as fertilisers. Name the salts which are produced by treating phosphate rock with: • sulfuric acid • phosphoric acid.	[2 marks]	
	Sulfuric acid Phosphoric acid		10

END OF QUESTIONS







Question number	Additional page, if required. Write the question numbers in the left-hand margin.



Question number	Additional page, if required. Write the question numbers in the left-hand margin.



Question number	Additional page, if required. Write the question numbers in the left-hand margin.



There are no questions printed on this page

DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED

Copyright information

For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from www.aqa.org.uk.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.

Copyright © 2023 AQA and its licensors. All rights reserved.



