Please check the examination details be	ring your candidate information	
Candidate surname		Other names
Centre Number Candidate N	umber	
Pearson Edexcel Leve	l 1/Lev	el 2 GCSE (9–1)
Tuesday 13 June 20	23	
Morning (Time: 1 hour 10 minutes)	Paper reference	1SC0/2CH
Combined Science PAPER 5	e	♦
		Higher Tier
You must have: Calculator, ruler		Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
 - there may be more space than you need.
- Calculators may be used.
- Any diagrams may NOT be accurately drawn, unless otherwise indicated.
- You must show all your working out with your answer clearly identified at the end of your solution.

Information

- The total mark for this paper is 60.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.
- In questions marked with an **asterisk** (*), marks will be awarded for your ability to structure your answer logically, showing how the points that you make are related or follow on from each other where appropriate.
- There is a periodic table on the back cover of the paper.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶



P72563A
©2023 Pearson Education Ltd.



Answer ALL questions. Write your answers in the spaces provided.

Some questions must be answered with a cross in a box \boxtimes . If you change your mind about an answer, put a line through the box \boxtimes and then mark your new answer with a cross \boxtimes .

A student used the apparatus shown in Figure 1 to investigate the reaction between marble chips and dilute hydrochloric acid.




Figure 1


The student recorded the volume of gas every minute as shown in Figure 2.

time in minutes	0 1		2	3	4	5	6	
volume of gas in cm ³	0	52	78	91	97	100	100	

Figure 2

(a) On the grid, plot the results shown in Figure 2.

Draw a curve of best fit.



(b) Rate of reaction can be calculated using

rate of reaction =
$$\frac{\text{volume of gas produced in 1 minute}}{\text{1 minute}}$$

Figure 3 shows the rates of reaction calculated from the results of this experiment.

The rate of reaction for the time interval 2 to 3 minutes is missing.

time interval	0 to 1	1 to 2	2 to 3	3 to 4	4 to 5	
	minute	minutes	minutes	minutes	minutes	
rate of reaction in cm³ min ⁻¹	52	26		6	3	

Figure 3

(i) Calculate the rate of reaction for the time interval 2 to 3 minutes.

(1)

rate of reaction =cm³ min⁻¹

(ii) State and explain what happens to the rate of reaction as the acid reacts with the marble chips in this experiment.

(3)

(c) The student repeated the experiment using the same volume of acid and the same mass of marble chips but used smaller marble chips.

All other conditions remained the same.

The student found that the reaction with the smaller marble chips was faster to start with but produced the same volume of gas.

Using this information, draw a line on the grid to show the results for the reaction with the smaller marble chips.

Label this line 'C'.

(2)

(Total for Question 1 = 9 marks)



2 Figure 4 shows some information about the group 1 metals.

group 1 metal	atomic number	relative atomic mass
lithium	3	7
sodium	11	23
potassium	19	39
rubidium	37	85
caesium	55	133

Figure 4

(a) Explain, in terms of their electronic configurations, why these metals are placed in group 1 of the periodic table.

(2)

(b) Which row shows two correct properties of group 1 metals?

(1)

- ⊠ A
- В
- \boxtimes D

properties of group 1 metals									
compounds are white in colour	high density								
low melting points	compounds are blue in colour								
soft enough to be cut by a knife	low melting points								
high density	conduct electricity								

(c)	The word equation for the reaction of potassium with bromine is	
(-)		
	potassium + bromine → potassium bromide	
	Add the missing state symbol and balance the equation for this reaction.	(2)
		(2)
	$K(Kgr(s)) + Br_2(g) \rightarrow Kgr(s)$	
	A sample of potassium contains three isotopes, potassium-39, potassium-40 and potassium-41.	
	(i) Explain the meaning of the term isotopes .	
		(2)
	(ii) This sample of potassium contains	
	93.25% potassium-39	
	0.02% potassium-40	
	6.73% potassium-41	
	Calculate the relative atomic mass of this sample of potassium.	
		(2)

relative atomic mass =

(Total for Question 2 = 9 marks)



(1)

(a) Figure 5 shows the percentage of three gases, **X**, **Y** and **Z**, in the Earth's early atmosphere.




Figure 5

What is the name of gas **Z**?

X argon

of gas in

- X carbon dioxide
- X C nitrogen
- X D oxygen



(b)	It is thought that small quantities of hydrogen sulfide, H₂S, were also in the Earth's	
()	early atmosphere.	
	Draw the dot and cross diagram for a molecule of hydrogen sulfide.	
	Show outer electrons only.	(2)
		(2)

(c) Acid rain is caused by some pollutant gases present in the atmosphere.

Explain how impurities in fossil fuels can result in acid rain.

(3)



(d) A student investigates the effect of acid rain on cress plants.

The student uses this method.

- step 1 grow 20 cress plants in each of two dishes, A and B
- **step 2** water the cress plants in dish **A** with 10 cm³ of dilute hydrochloric acid with a pH of 2
- step 3 water the cress plants in dish B with 10 cm³ of pure water with a pH of 7
- **step 4** repeat steps 2 and 3 every day for one week
- **step 5** count how many plants are still alive after one week.
- (i) State what piece of equipment the student could use to measure the pH of each liquid.

(1)

(ii) Explain **one** improvement that the student could make to the method to make the results more valid.

(2)

(Total for Question 3 = 9 marks)

8





BLANK PAGE



4 Chlorine gas can be prepared by reacting concentrated hydrochloric acid with solid potassium manganate(VII).

Figure 6 shows the apparatus used.




Figure 6

(a) Figure 7 shows the hazard symbols for concentrated hydrochloric acid, potassium manganate(VII) and chlorine gas.

substance	hazard symbol
concentrated hydrochloric acid	
potassium manganate(VII)	
chlorine gas	

Figure 7

Use the information in Figure 7 to help you answer (a)(i) and (a)(ii).



(Total for Question 4 = 9	marks)									
(d) In the reaction, potassium manganate(VII), KMnO4, reacts with hydrochloric aci to form manganese chloride, MnCl2, potassium chloride, chlorine and water. Write the balanced equation for the reaction.	id (3)									
(c) Suggest why damp blue litmus is placed at the top of the gas jar.										
(b) State the purpose of the delivery tube.										
reason										
(ii) Explain one precaution that should be taken when preparing the sample of chlorine gas.precaution										
 C oxidising, harmful and hazardous to the environment D oxidising, toxic and corrosive 										
A flammable, harmful and corrosiveB flammable, toxic and hazardous to the environment										
(i) What are the hazards associated with potassium manganate(VII)?										



- **5** Ammonia can be produced from the reaction of hydrogen with nitrogen.
 - (a) What is the percentage by mass of nitrogen in ammonia, NH_3 ?

(relative atomic masses: H = 1.0, N = 14)

(1)

- A 18%
- **■ B** 42%
- **C** 51%
- **D** 82%
- (b) The reaction between hydrogen and nitrogen is exothermic.

Figure 8 shows the reaction profile of this exothermic reaction.




Figure 8

(i) Which arrow represents the activation energy for the reaction?

(1)

- A arrow P
- B arrow Q
- C arrow R
- **D** arrow **S**

(ii)	Describe what the reaction profile shows about the energy involved in
	bond breaking and bond making in this reaction.

(2)

(iii) Figure 9 shows the energies of some bonds.

bond	bond energy in kJ mol ⁻¹
N≡N	944
Н—Н	436
H—N	388

Figure 9

The equation for the reaction between nitrogen and hydrogen to form ammonia is

$$N \equiv N + 3 H - H \rightarrow 2 \begin{array}{c} H \\ N \end{array}$$

Calculate the energy change, in kJ mol⁻¹, for this reaction.

(4)

|
 |
|------|------|------|------|------|------|------|------|------|------|
|
 |
|
 |
|
 |
| | | | | | | | | | |
| | | | | | | | | | |

energy change =kJ mol⁻¹



(Total for Question 5 = 11 marks)

(c)	Ammonia, $\mathrm{NH_3}$, and silicon dioxide, $\mathrm{SiO_2}$, are both compounds that are made of two non-metallic elements.	
	Ammonia has a boiling point of –33 °C. Silicon dioxide has a boiling point of 2230 °C.	
	Explain why the boiling points of ammonia and silicon dioxide are so different.	
	, , , , , , , , , , , , , , , , , , , ,	(3)


BLANK PAGE



6 Crude oil is a mixture of hydrocarbons.

Crude oil can be separated into useful fractions by the process of fractional distillation in a fractionating column.

(a) Figure 10 shows a fractionating column, the fractions obtained and the trend in viscosity of the fractions.



(i) Which row shows the correct uses for bitumen, diesel oil and fuel oil?

(1)

		bitumen	diesel oil	fuel oil
×	A	fuel for large ships	surfacing roads	fuel for trains
X	В	fuel for large ships	fuel for trains	surfacing roads
X	C	surfacing roads	fuel for trains	fuel for large ships
X	D	surfacing roads	fuel for large ships	fuel for trains

(ii)	Explain the trend in the viscosity of the fractions.	
------	--	--

-//	\neg	٠,
	- 11	-1

(b) Hydrocarbon \mathbf{X} was cracked to form one molecule of hexane, C_6H_{14} , and one molecule of alkene \mathbf{Y} .

$$\boldsymbol{X} \ \rightarrow \ C_6 H_{14} \ + \ \boldsymbol{Y}$$

The relative formula mass of Y is 56.

The empirical formula of \mathbf{Y} is CH_2 .

Deduce the molecular formula of hydrocarbon X.

Show your working.

(relative atomic masses: H = 1.0, C = 12)

(4)

molecular formula of **X** =



(6)

*(c) Large quantities of methane are used as a fuel.

Figure 11 shows a Bunsen burner.

Methane can be used as fuel for the Bunsen burner.




Figure 11

The air-hole on the chimney of the Bunsen burner can be opened and closed.

Explain the effect of opening and closing the air-hole of the Bunsen burner on the products of combustion of methane and the harm that using large quantities of methane as a fuel can cause.

 	 •••••	•••••	

(Total for Question 6 = 13 marks)
(
TOTAL FOR PAPER = 60 MARKS
I O IAL FOR FAFER - UU IVIARRO



The periodic table of the elements

0	4 He helium 2	20 Ne neon 10	40 Ar argon 18	84 Kr krypton 36	131 Xe xenon 54	[222] Rn radon 86
7		19 F fluorine 9	35.5 CI chlorine 17	80 Br bromine 35	127 	[210] At astatine 85
9		16 O	32 S sulfur 16	79 Se selenium 34	128 Te tellurium 52	[209] Po polonium 84
2		14 N nitrogen 7	31 P phosphorus 15	75 As arsenic 33	122 Sb antimony 51	209 Bi bismuth 83
4		12 C carbon 6	28 Silicon 14	73 Ge germanium 32	119 Sn th	207 Pb lead lead 82
က		11 B boron 5	27 AI aluminium 13	70 Ga gallium 31	115 In indium 49	204 T thallium 81
	•			65 Zn zinc 30	112 Cd cadmium 48	201 Hg mercury 80
				63.5 Cu copper 29	108 Ag silver 47	197 Au gold 79
				59 Ni nickel 28	106 Pd palladium 46	195 Pt platinum 78
				59 Co cobalt 27	103 Rh rhodium 45	192 Ir iridium 77
	1 X hydrogen				Ru ruthenium 44	190 Os osmium 76
•				55 Mn manganese 25	[98] Tc technetium 43	186 Re rhenium 75
	Key	nass ool umber		52 Cr chromium 24	96 Mo molybdenum 42	184 W tungsten 74
		relative atomic mass atomic symbol name atomic (proton) number		51 V vanadium 23	93 Nb niobium 41	181 Ta tantalum 73
		relati atc atomic		48 Ti titanium 22	91 Zr zirconium 40	178 Hf hafnium 72
				45 Sc scandium 21	89 Y yttrium 39	139 La * lanthanum 57
2		9 Be beryllium 4	24 Mg magnesium 12	40 Ca calcium 20	88 Sr strontium 38	137 Ba barium 56
_		7 Li lithium 3	23 Na sodium 11	39 K potassium 19	85 Rb rubidium 37	133 Cs caesium 55

^{*} The elements with atomic numbers from 58 to 71 are omitted from this part of the periodic table.

The relative atomic masses of copper and chlorine have not been rounded to the nearest whole number.