Please check the examination deta	ils bel	ow before ente	ring your candidate information
Candidate surname			Other names
Pearson Edexcel Level 1/Level 2 GCSE (9–1)	Cen	tre Number	Candidate Number
<b>Time</b> 1 hour 10 minutes		Paper reference	1SC0/2CF
Combined Scient PAPER 5 Foundation Tier	nc	e	
You must have: Calculator, ruler			Total Marks

# **Instructions**

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions.
- Answer the questions in the spaces provided
  - there may be more space than you need.
- Calculators may be used.
- Any diagrams may NOT be accurately drawn, unless otherwise indicated.
- You must show all your working out with your answer clearly identified at the end of your solution.

## Information

- The total mark for this paper is 60.
- The marks for each question are shown in brackets
  - use this as a guide as to how much time to spend on each question.
- In questions marked with an **asterisk** (\*), marks will be awarded for your ability to structure your answer logically showing how the points that you make are related or follow on from each other where appropriate.
- There is a periodic table on the back cover of the paper.

### **Advice**

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- Good luck with your examination.

Turn over ▶







# Answer ALL questions. Write your answers in the spaces provided.

Some questions must be answered with a cross in a box  $\boxtimes$ . If you change your mind about an answer, put a line through the box  $\boxtimes$  and then mark your new answer with a cross  $\boxtimes$ .

1	Magnesium reacts with	dilute sulfuric acid to	form magnesium	sulfate and hydrogen gas.
	9			, , ,

A student wants to find out the effect of temperature on the rate of this reaction.

The student used the following method.

- step 1 pour 25 cm³ of dilute sulfuric acid into a conical flask
- step 2 warm the acid until its temperature is 30 °C
- step 3 add a piece of magnesium to the acid
- **step 4** start a stopwatch
- **step 5** wait until the reaction has finished
- **step 6** stop the stopwatch
- step 7 repeat steps 1-6 but at 50°C.

**D** thermometer

(a) The student kept the volume of sulfuric acid the same when they repeated the method at 50 °C.

State two other variables that should be kept the same.

1			
2			
	(b) Which	n pie	cce of equipment can be used to find the volume of 25 cm <sup>3</sup> of sulfuric acid?
	X	A	balance
	$\times$	В	measuring cylinder
	$\times$	C	ruler

(c) State how the student will know that the reaction has finished.

(1)

(2)

(d) The reaction at  $50\,^{\circ}\text{C}$  was faster than the reaction at  $30\,^{\circ}\text{C}$ .

Give **one** reason, in terms of particles, why the reaction at  $50^{\circ}$ C was faster than the reaction at  $30^{\circ}$ C.

(1)



X

average rate of reaction =	cm³ s <sup>-1</sup>
reaction.	(2)
Calculate the average rate of reaction, in cm <sup>3</sup> s <sup>-1</sup> , for the first 60 seconds of the	
(e) At 50 °C, 15.0 cm <sup>3</sup> of gas was produced during the first 60 seconds of the reaction.	

- **2** This question is about the noble gases.
  - (a) (i) State, in terms of outer shell electrons, why the noble gases are unreactive.

(1)

(ii) Figure 1 shows an airship, filled with helium, floating above the ground.




Figure 1

Helium, hydrogen and krypton are gases. Figure 2 shows the reactivity and density, at room temperature and pressure, of helium, hydrogen and krypton.

gas	reactivity	density in g cm <sup>-3</sup>
helium	unreactive	0.00018
hydrogen	very reactive	0.00009
krypton	unreactive	0.00380

Figure 2

The density of air is  $0.001225 \,\mathrm{g}\,\mathrm{cm}^{-3}$ .

Helium is used in airships.

Explain why hydrogen and why krypton are **not** used in airships.

//	2	1
ı	-5	)
-1	$\sim$	11


(b) Mendeleev produced one of the earliest periodic tables.

State why he could **not** include any of the noble gases in his periodic table.

(1)


(c) Figure 3 shows the boiling points of some of the noble gases.

noble gas	boiling point in °C
helium	-269
neon	-246
argon	-186
krypton	-153
xenon	

Figure 3

(i) Complete the bar chart to show the boiling points of neon and argon.





(ii) Predict the boiling point of xenon.

(1)

boiling point of xenon = .....°C

(Total for Question 2 = 8 marks)




3	Fluorine, chlorine, bromine and iodine are elements in group 7 of the periodic table.  (a) (i) State the name given to the group 7 elements.								
			e other element that is in group 7. periodic table on the back of this exam pa	aper to help you.	(1)				
	(iii) Whi	ch el	ement is liquid at room temperature and	pressure?					
	$\times$	Α	fluorine		(1)				
		В	chlorine						
	$\boxtimes$	C	bromine						
	×	D	iodine						
	(iv) Whi	ch el	ement is dark grey in colour at room tem	perature and pressure?	(4)				
		Α	fluorine		(1)				
	$\boxtimes$	В	chlorine						
	$\boxtimes$	C	bromine						
	$\boxtimes$	D	iodine						
			th chlorine to form tin chloride. tin chloride contains 1.19 g of tin and 1.42	2 g of chlorine.					
	Calculat	te th	e empirical formula of this tin chloride.						
	(relative	ato	nic masses: Cl = 35.5, Sn = 119.0)						
	You mu	st sh	ow your working.		(3)				
•••••									
•••••			empirical formula of	this tin chloride =					
			- p						



(c) Tin also reacts with fluorine.

The reaction between fluorine and tin is much more vigorous than the reaction between chlorine and tin.

Figure 4 shows the electronic configurations of fluorine and chlorine.






Figure 4


Explain, in terms of their electronic configurations, why fluorine reacts with tin more vigorously than chlorine reacts with tin.

(2)

(Total for Question 3 = 9 marks)



# **BLANK PAGE**






Figure 5

(a) (i) Write the word equation for hydrogen peroxide solution decomposing.

(1)


(ii) In this reaction hydrogen peroxide is a solution, water is a liquid and oxygen is a gas.


Draw one straight line from each substance to its correct state symbol.

(2)


## substance

hydrogen peroxide solution






# state symbol







(b) Describe the test to show the gas produced is oxygen.

(2)



(2)

(c) Figure 6 shows the electron arrangement for an atom of hydrogen and an atom of oxygen.




Figure 6

Complete the dot and cross diagram in Figure 7 for a molecule of water,  $\rm H_2O$ . Draw outer shell electrons only.




Figure 7

(d) Liver contains the enzyme catalase.

A piece of liver was added to another sample of hydrogen peroxide solution. Figure 8 shows the results.





Figure 8





**5** (a) The concentration of a solution can be calculated using the equation

concentration of solution = 
$$\frac{\text{mass of solid}}{\text{volume of solution}}$$


A student dissolved 9.25 g of ammonium chloride in water and made up the solution to a volume of 200 cm<sup>3</sup>.

Use the equation to calculate the concentration of this solution in g dm<sup>-3</sup>.

(2)

concentration = ...... g dm<sup>-3</sup>

(b) Dissolving ammonium chloride in water is an endothermic process. Figure 10 shows part of the reaction profile for this process.




progress of reaction

Figure 10

(i) Explain how Figure 10 shows that dissolving ammonium chloride in water is an endothermic process.


(2)



(ii) Complete the reaction profile in Figure 10 and label the activation energy.

(2)

(c) A student used the equipment in Figure 11 to investigate whether electricity can pass through solid ammonium chloride and through ammonium chloride solution.






Figure 11

If an electrical current flows in the circuit, the lamp will light up.

Figure 12 shows the results of the investigation.

substance	lamp
solid ammonium chloride	did not light up
ammonium chloride solution	lit up brightly

Figure 12

Explain the results of the investigation.


- (d) Ammonia gas is toxic.
  - (i) Which symbol should be placed on a container of a toxic gas?

(1)











(ii) Give **one** safety precaution that should be taken when working with toxic gases in the laboratory.

(1)

(Total for Question 5 = 11 marks)

	Stat	e th	e name of the process used to separate diesel oil from crude oil.	(1)		
(b) Diesel oil contains alkanes. These alkanes are part of an homologous series.						
	Which statement about compounds in this homologous series is true?					
	×	A	they have the same chemical formula	(1)		
	×	В	they have the same empirical formula			
	X	C	they have the same general formula			
	X	D	they have the same molecular formula			
(c) When fuels such as diesel oil are burned, the high temperatures produced can cause nitrogen and oxygen in the air to form the pollutant nitrogen dioxide. Complete the balanced equation for the reaction.						
(d)			$N_2 + 2O_2 \rightarrow$ how the greenhouse effect is caused by the gases produced by the te combustion of diesel oil.	(3)		

*(e)	Diesel oil can contain impurities of sulfur.  Burning diesel oil containing impurities of sulfur can result in acid rain.  Acid rain is harmful to the environment.						
	Explain how acid rain is formed and the harm it can do.	(6)					
		(6)					

	(Total for Question 6 = 13 marks)		
	(		
TOTAL FOR PAPER = 60 MARKS			



**BLANK PAGE** 

# **BLANK PAGE**



# The periodic table of the elements

0	4 <b>He</b> helium 2	20 <b>Ne</b> neon 10	40 <b>Ar</b> argon 18	84 <b>Kr</b> krypton 36	131 <b>Xe</b> xenon 54	[222] <b>Rn</b> radon 86
7		19 <b>F</b> fluorine 9	35.5 <b>CI</b> chlorine 17	80 <b>Br</b> bromine 35	127 	[210] <b>At</b> astatine 85
9		16 <b>O</b> oxygen 8	32 <b>S</b> sulfur 16	79 <b>Se</b> selenium 34	128 <b>Te</b> tellurium 52	[209] <b>Po</b> Polonium 84
2		14 N nitrogen 7	31 P phosphorus 15	75 <b>As</b> arsenic 33	122 <b>Sb</b> antimony 51	209 <b>Bi</b> bismuth 83
4		12 <b>C</b> carbon 6	28 <b>Si</b> silicon 14	73 <b>Ge</b> germanium 32	119 <b>Sn</b> tin 50	207 <b>Pb</b> lead 82
က		11 <b>B</b> boron 5	27 AI aluminium 13	70 <b>Ga</b> gallium 31	115 In indium 49	204 <b>TI</b> thallium 81
	'			65 <b>Zn</b> zinc 30	112 <b>Cd</b> cadmium 48	201 <b>Hg</b> mercury 80
				63.5 <b>Cu</b> copper 29	108 <b>Ag</b> silver 47	197 <b>Au</b> gold 79
				59 <b>Ni</b> nickel 28	106 <b>Pd</b> palladium 46	195 <b>Pt</b> platinum 78
				59 <b>Co</b> cobalt 27	103 <b>Rh</b> rhodium 45	192 <b>Ir</b> iridium 77
	1 Hydrogen 1			56 iron 26	101 <b>Ru</b> ruthenium 44	190 <b>Os</b> osmium 76
!				55 Mn manganese 25	[98] Tc technetium 43	186 <b>Re</b> rhenium 75
		relative atomic mass  atomic symbol  atomic (proton) number		52 Cr	96 <b>Mo</b> molybdenum 42	184 <b>W</b> tungsten 74
	Key			51 V vanadium 23	93 <b>Nb</b> niobium 41	181 <b>Ta</b> tantalum 73
				48 Ti titanium 22	91 Zr zirconium 40	178 <b>Hf</b> hafnium 72
				45 Sc scandium 21	89 <b>Y</b> 39	139 <b>La*</b> Ianthanum 57
2		9 <b>Be</b> beryllium 4	24 <b>Mg</b> magnesium 12	40 <b>Ca</b> calcium 20	88 Sr strontium 38	137 <b>Ba</b> barium 56
_		7 <b>Li</b> Ilthium 3	23 <b>Na</b> sodium 11	39 <b>K</b> potassium 19	85 <b>Rb</b> rubidium 37	133 <b>Cs</b> caesium 55

<sup>\*</sup> The elements with atomic numbers from 58 to 71 are omitted from this part of the periodic table.

The relative atomic masses of copper and chlorine have not been rounded to the nearest whole number.