

Please write clearly in block capitals.	
Centre number	Candidate number
Surname	
Forename(s)	
Candidate signature	

AS CHEMISTRY

Paper 1 Inorganic and Physical Chemistry

Tuesday 22 May 2018

Morning

Time allowed: 1 hour 30 minutes

Materials

For this paper you must have:

- the Periodic Table/Data Sheet, provided as an insert (enclosed)
- a ruler with millimetre measurements
- a scientific calculator, which you are expected to use where appropriate.

Instructions


- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer all questions.
- You must answer the questions in the spaces provided. Do **not** write outside the box around each page or on blank pages.
- All working must be shown.
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80.

Advice

• You are advised to spend about 65 minutes on Section A and 25 minutes on Section B.





Section A


Answer all questions in this section.

0 1 This question is about atomic structure.

> In the nineteenth century JJ Thomson discovered the electron. He suggested that negative electrons were found throughout an atom like 'plums in a pudding of positive charge'.

Figure 1 shows an atom of element R using the 'plum pudding' model. An atom of **R** contains seven electrons.

Figure 1



0	1	.[1	State two differences between the 'plum pudding' model and the model of atomic
				structure used today.

[2 marks]

1			

0 1 . 2	Deduce the full electron configuration of an atom of element R .	
		[1 mark]

0 1 . 3	Identify R and deduce the formula of the compound formed when R reacts with the
	Group 2 metal in the same period as R .
	[1 mar

[1 mark]



	3
0 2	This question is about sodium fluoride (NaF).
	Some toothpastes contain sodium fluoride. The concentration of sodium fluoride can be expressed in parts per million (ppm). 1 ppm represents a concentration of 1 mg in every 1 kg of toothpaste.
0 2.1	A 1.00 g sample of toothpaste was found to contain 2.88 x 10^{-5} mol of sodium fluoride.
	Calculate the concentration of sodium fluoride, in ppm, for the sample of toothpaste. Give your answer to 3 significant figures.
	[4 marks]

Concentration of sodium fluoride



ppm



0 2 . 2	Sodium fluoride is toxic in high concentrations. Major health problems can occur if concentrations of sodium fluoride are greater than 3.19×10^{-2} g per kilogram of body mass. Deduce the maximum mass of sodium fluoride, in mg, that a 75.0 kg person could swallow without reaching the toxic concentration. [1 marks]	
	Mass of sodium fluoridemg	
0 2.3	The concentration of sodium fluoride in a prescription toothpaste is 2800 ppm. Use your answer to Question 02.2 to deduce the mass of toothpaste, in kg, that a 75.0 kg person could swallow without reaching the toxic concentration. [1 mar	k]
	Mass of toothpastekg	



0 2 . 4 Identify the diagram in **Figure 2** that shows the correct relative sizes of the ions in sodium fluoride.

Justify your answer.

[3 marks]

_






Diagram	 =		
Justification			

Turn over for the next question



A student heated a solid sample of Na_2CO_3 . xH_2O for 1 minute to remove water and determine a value for x

Figure 3 shows the apparatus used. Table 1 shows the results recorded.

Figure 3




Table 1

Mass of empty evaporating basin	24.35 g
Mass of evaporating basin and solid before heating	25.47 g
Mass of evaporating basin and solid after heating for 1 minute	24.92 g

Use the data in **Table 1** to calculate a value for x in the formula Na₂CO₃. xH₂O Give your answer to 2 decimal places.

[5 marks]



0 3.2	The correct value for <i>x</i> is 10
	Suggest a reason for the difference between the experimental value for x and the correct value. (If you were unable to calculate an experimental value for x assume it was 8.05. This is not the correct experimental value.)
0 3.3	Suggest how the procedure could be improved, using the same apparatus, to give a more accurate value for x Justify your answer.
	Suggestion
	Justification
	Justification

Turn over for the next question

Turn over ▶



0 4.1	Separate unlabelled solid samples of three anhydrous sodium compounds are provided for a student to identify. These compounds are known to be sodium carbonate, sodium fluoride and sodium chloride but it is not known which sample is which. Outline a logical sequence of test-tube reactions that the student could carry out to identify each of these compounds. Include the observations the student would expect to make. Give equations, including state symbols, for any reactions that would take place. [6 marks]



Town areas for the most arrestion
Turn over for the next question



This question is about equilibrium.


Sulfur trioxide decomposes to form sulfur dioxide and oxygen at temperature T_1 according to the equilibrium shown.

$$2SO_3(g) \rightleftharpoons 2SO_2(g) + O_2(g)$$

$$\Delta H = +196 \text{ kJ mol}^{-1}$$

The graph in **Figure 4** shows the concentrations of sulfur trioxide and of oxygen over a period of 6 minutes at temperature T_1

Figure 4



0 5 . 1

State the time, to the nearest minute, when equilibrium is first established. Explain your answer.

[2 marks]

Time	minutes
	•

Explanation

0 5.2	Sketch on the graph in Figure 4 how the concentration of sulfur dioxide changes over these 6 minutes at temperature <i>T</i> ₁ [2 marks]
0 5 . 3	The temperature of the mixture was changed to T_2 and the mixture left to establish a new equilibrium. In the new equilibrium mixture the concentration of sulfur trioxide was found to be 0.07 mol dm^{-3} Deduce which of T_1 and T_2 is the higher temperature. Explain your deduction. [2 marks] Higher temperature Explanation

Turn over for the next question



A student determined the relative molecular mass, M_r , of an unknown volatile liquid **Y** in an experiment as shown in **Figure 5**.

The student used a hypodermic syringe to inject a sample of liquid ${\bf Y}$ into a gas syringe in an oven.

At the temperature of the oven, liquid Y vaporised.

The student's results are shown in Table 2.

Figure 5




Table 2

Mass of hypodermic syringe and liquid Y before injection	10.91 g
Mass of hypodermic syringe and liquid Y after injection	10.70 g
Oven temperature	98.1 °C
Atmospheric pressure	102 kPa
Increase in volume in gas syringe after injection of Y	85.0 cm ³



#
找名校职
₹
íń.
뿗
₻
픨
壍
草线山
₩ —
捶
中難力
4(1)
企
渡信
!!!!
忈
程序
回名
Ш
_

0 6.1	Define the term relative molecular mass (M_r) .
	Use the experimental results in Table 2 to determine the relative molecular mass of Y . The gas constant $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ [5 marks]
0 6 . 2	Some of the liquid injected did not evaporate because it dripped into the gas syringe nozzle outside the oven.
	Explain how this would affect the value of the $M_{\rm r}$ of Y calculated from the experimental results. [2 marks]





0 7	Chlorine is used to decrease the numbers of microorganisms in water.
	When chlorine is added to water, there is a redox reaction, as shown by the equation
	$Cl_2 + H_2O \Rightarrow HClO + HCl$
0 7.1	Deduce the oxidation state of chlorine in HClO and the oxidation state of chlorine in HCl
	[1 mark]
	Oxidation state of chlorine in HClO
	Oxidation state of chlorine in HCl
0 7.2	Give two half-equations to show the oxidation and reduction processes that occur in this redox reaction. [2 marks]
	Oxidation half-equation
	Reduction half-equation
0 7.3	Chlorine is reacted with cold, aqueous sodium hydroxide in the manufacture of bleach. Give an equation for this reaction between chlorine and sodium hydroxide.
	[1 mark]



0 7 .

The concentration of ClO⁻ ions in bleach solution can be found by reaction with iodide ions.

The overall equation for this reaction is shown.

$$\text{ClO}^- \ + \ 2\text{I}^- \ + \ 2\text{H}^+ \ \rightarrow \ \text{I}_2 \ + \ \text{Cl}^- \ + \ \text{H}_2\text{O}$$

A sample of bleach solution was found to contain ClO-ions with a concentration of 0.0109 mol dm⁻³

Potassium iodide is added to a 20.0 cm³ portion of this bleach solution.

Calculate the mass, in mg, of potassium iodide needed to react with all of the ClO⁻ions in the sample of bleach.

Give your answer to the appropriate number of significant figures.

Give **one** observation during this reaction.

[4 marks]

	Mass of potassium iodide	mg
Observation		

Question 7 continues on the next page





0 7 . 5 Potassium chlorate(VII), KClO₄, is used in fireworks. When potassium chlorate(VII) decomposes, it produces potassium chloride and oxygen.

Give an equation for the decomposition of potassium chlorate(VII). Use the data in **Table 3** to calculate the enthalpy change for this reaction.

[2 marks]

Table 3

Substance	$\Delta_{\mathrm{f}} H$ / kJ mol ⁻¹
KClO ₄ (s)	-434
KCl(s)	-436

Equation			

Enthalpy change	kJ mol ⁻¹



[5 marks]

0 8 A sample of bromine was analysed in a time of flight (TOF) mass spectrometer and found to contain two isotopes, ⁷⁹Br and ⁸¹Br

After electron impact ionisation, all of the ions were accelerated to the same kinetic energy (KE) and then travelled through a flight tube that was 0.950 m long.

0 8 . **1** The ⁷⁹Br⁺ ions took 6.69 x 10⁻⁴ s to travel through the flight tube.

Calculate the mass, in kg, of one ion of ⁷⁹Br⁺ Calculate the time taken for the ⁸¹Br⁺ ions to travel through the same flight tube.

The Avogadro constant, $L = 6.022 \times 10^{23} \text{ mol}^{-1}$

$$KE = \frac{1}{2} mv^2$$
 where $m = \text{mass (kg)}$ and $v = \text{speed (m s}^{-1})$

$$v = \frac{d}{t}$$
 where $d = \text{distance (m)}$ and $t = \text{time (s)}$

Mass of one ion of ⁷⁹Br⁺ kg

Time taken by ⁸¹Br⁺ ions

Turn over ▶



	18
0 8.2	Explain how ions are detected and relative abundance is measured in a TOF mass spectrometer. [2 marks]




0 9	This question is about compounds containing fluorine.	
0 9 . 1	Draw the shape of a molecule of krypton difluoride (KrF $_2$). Include in your answer any lone pairs of electrons that influence the shape. Name the shape produced by the atoms in a KrF $_2$ molecule and suggest a bond angle.	[3 marks]
	Name of shape Bond angle	
0 9.2	There are two lone pairs of electrons on the oxygen atom in a molecule of oxygen difluoride (OF ₂). Explain how the lone pairs of electrons on the oxygen atom influence the boin oxygen difluoride.	nd angle [2 marks]



0 9 . 3	Silicon tetrafluoride (SiF ₄) is a tetrahedral molecule.
	Deduce the type of intermolecular forces in SiF ₄ Explain how this type of intermolecular force arises and why no other type of intermolecular force exists in a sample of SiF ₄ [3 marks]
	Intermolecular forces in SiF ₄
	Explanation









找名校导师,用小草线上辅导(微信小程序同名)

Section B

Answer all questions in this section.

Only **one** answer per question is allowed.

For each answer completely fill in the circle alongside the appropriate answer.

CORRECT METHOD

WRONG METHODS

∞ •	(
-----	---

If you want to change your answer you must cross out your original answer as shown.



If you wish to return to an answer previously crossed out, ring the answer you now wish to select as shown. 🏲

You may do your working in the blank space around each question but this will not be marked. Do **not** use additional sheets for this working.

0 Which row shows the bonding in ammonium chloride?

[1 mark]

	Covalent	Dative covalent	lonic	
A	✓	*	*	0
В	✓	✓	*	0
С	✓	✓	✓	0
D	*	×	✓	0

1 1 How many protons are there in 6.0 g of nitrogen gas?

Avogadro constant, $L = 6.022 \times 10^{23} \text{ mol}^{-1}$

[1 mark]


A 1.3×10^{23}

B 9.0×10^{23}

 $\mathbf{C} \ 1.8 \times 10^{24}$

D 3.6×10^{24}





What is the property?

[1 mark]

- A Atomic radius
- 0
- **B** Electronegativity
- 0
- C First ionisation energy
- 0

D Melting point

- 0
- 1 3 A 30 cm³ sample of nitrogen was reacted with a 60 cm³ sample of fluorine according to the equation

$$\frac{1}{2} N_2(g) + \ \frac{3}{2} F_2(g) \ \to \ NF_3(g)$$

What is the volume of the gas mixture after the reaction, at constant temperature and pressure?

[1 mark]

A 20 cm³

0

B 30 cm³

0

C 40 cm³

0

D 50 cm³



1 4	Which substance is used to reduce titanium(IV) chloride in the extraction of titaniu metal?		
		[1 mar	«]
	A Magnesium		
	B Manganese		
	C Vanadium		
	D Zinc		
1 5	Which statement about barium sulfate is correct?		
		[1 mar	(]
	A It is soluble in water at a temperature of 100 °C.	0	
	B It is used in medicine because it does not dissolve in body fluids.	0	
	C It is a pale yellow solid.	0	
	D It reacts with acidified barium chloride solution.	0	
1 6	Which statement is correct about the reaction between concentrated s	ulfuric acid and	
	solid sodium bromide?	[1 marl	k]
	A Bromide ions are reduced.	0	
	B Hydrogen bromide and sulfur are formed.		
	C Sulfuric acid acts as an oxidising agent.		
	D Bromine and hydrogen sulfide are formed.		
	Distinic and Hydrogen sumde are formed.		



1 7	Which compound is used to treat the symptoms of indigestion?		[1 mark]
	A MgO	0	
	B Mg(OH) ₂	0	
	C CaO	0	
	D Ca(OH) ₂	0	
1 8	Which element has the highest fi	rst ionisation energy?	[1 mark]
	A Aluminium		
	B Phosphorus		
	C Silicon	0	
	D Sulfur	0	
1 9	A solution of volume 500 cm ³ cor	ntains 150 g of ammonia.	
	What is the concentration, in mol	dm ⁻³ , of ammonia in this solution?	[1 mark]
	A 0.54		
	A 0.51		
	B 8.82	0	
	C 16.7		
	D 17.6	0	



Refer to the following information when answering Questions 20, 21, 22, 23 and 24.

A student devised an experiment to find the concentration of sulfuric acid in a sample of battery acid.


- A measuring cylinder was used to transfer 10 cm³ of battery acid to a volumetric flask.
- Distilled water was added to the volumetric flask until the volume reached 250 cm³
- A 25.0 cm³ sample of diluted acid was transferred from the volumetric flask to a conical flask using a pipette.
- A few drops of methyl orange indicator were added to the acid in the conical flask before titrating the acid with sodium hydroxide.
- The titration was repeated five times but concordant results were **not** obtained. (Note: Methyl orange is red in acid and yellow in alkali.)

Which suggestion would improve the chances of obtaining concordant titres?	[1 mark]
A Invert the volumetric flask several times after adding the distilled water.	
B Wash the pipette with distilled water between each titration.	
C Add extra drops of indicator to the sample when nearing the end point in each titration.	
D Use a more concentrated solution of sodium hydroxide in the burette.	
Which suggestion about rinsing the conical flask between each titration would imaccuracy of the titrations?	prove the
A Rinsing with acid.	
B Rinsing with alkali.	
C Rinsing with water.	
D No rinsing with any liquid.	
	A Invert the volumetric flask several times after adding the distilled water. B Wash the pipette with distilled water between each titration. C Add extra drops of indicator to the sample when nearing the end point in each titration. D Use a more concentrated solution of sodium hydroxide in the burette. Which suggestion about rinsing the conical flask between each titration would imaccuracy of the titrations? A Rinsing with acid. B Rinsing with alkali. C Rinsing with water.



2 2	Which suggestion would reduce the overall measurement uncertainty in	n the titrati	on? [1 mark]
	A Use less concentrated alkali in the burette.	0	
	B Use phenolphthalein indicator instead of methyl orange.	0	
	C Use smaller samples of the diluted acid in each titration.	0	
	D Begin each titration with the burette filled to the 0.00 cm ³ mark.	0	
		f?	
2 3	Which of these is important in ensuring that the student's experiment is	sare?	[1 mark]
	A Do the titration in a fume cupboard.	0	
	B Wear gloves when measuring out the battery acid.	0	
	C Wash hands before doing the titration.	0	
	D Carry the burette horizontally when collecting the apparatus.	0	
2 4	Which colour change is observed at the end point in each titration?		[1 mark]
	A Yellow to red		
	B Red to orange		
	C Yellow to orange		
	D Red to yellow		
	END OF QUESTIONS		





Copyright information

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2018 AQA and its licensors. All rights reserved.

