

| Please write cle | early in block capitals. |
|------------------|--|
| Centre number | Candidate number |
| Surname | |
| Forename(s) | |
| Candidate sign | lature I declare this is my own work. |
| | i deciale tills is tily own work. |

AS CHEMISTRY

Paper 1 Inorganic and Physical Chemistry

Tuesday 16 May 2023

Morning

Time allowed: 1 hour 30 minutes

Materials

For this paper you must have:

- the Periodic Table/Data Sheet, provided as an insert (enclosed)
- a ruler with millimetre measurements
- a scientific calculator, which you are expected to use where appropriate.

Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer all questions.
- You must answer the questions in the spaces provided. Do **not** write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- All working must be shown.
- Do all rough work in this book. Cross through any work you do not want to be marked.

| For Examiner's Use | | | |
|--------------------|------|--|--|
| Question | Mark | | |
| 1 | | | |
| 2 | | | |
| 3 | | | |
| 4 | | | |
| 5 | | | |
| 6 | | | |
| 7 | | | |
| 8 | | | |
| Section B | | | |
| TOTAL | | | |

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80.

Advice

You are advised to spend about 65 minutes on Section A and 25 minutes on Section B.



Section A

Answer all questions in this section.

| 0 | 1 | This question is about the elements in Period 3. |
|---|---|--|
|---|---|--|

0 1. Give the full electron configuration of the element in Period 3 with the highest first ionisation energy.

[1 mark]

0 1. 2 Give an equation, including state symbols, to represent the process that occurs when the second ionisation energy of sodium is measured.

[1 mark]

0 1 . 3 Table 1 shows some successive ionisation energies for an element in Period 3.

Table 1

| Ionisation number | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
|--|------|------|------|------|------|------|--------|-------|
| lonisation energy / kJ mol ⁻¹ | 1000 | 2260 | 3390 | 4540 | 6990 | 8490 | 27 100 | 31700 |

Element

Identify the Period 3 element.

Explain your answer.

[3 marks]

| Explanation | | | |
|-------------|--|--|--|



| 0 2 . 1 | This question is about the elements in Group 2. Describe the structure and bonding in magnesium. [2 marks] |
|---------|---|
| 0 2.2 | State the trend in the atomic radius of the elements down Group 2 from Mg to Ba Give a reason for this trend. |
| | Trend |
| 0 2.3 | Give an equation, including state symbols, for the reaction of magnesium with steam. State two observations for this reaction. [3 marks] |
| | Observation 1 Observation 2 |
| | Question 2 continues on the next page |





| Find Personal Tutor from www.wisesprout.co.uk |
|---|
| 找名校导师, |
| ,用小草线上辅导 |
| (微信小程序同名) |

| 0 2 . 4 | The sulfates of the elements in Group 2 from Mg to Ba have different solubilities. |
|---------|--|
| | State the formula of the least soluble of these sulfates. |
| | Give a use for this sulfate. [2 marks] |
| | Formula |
| | Use |
| | |

0 2 . 5 A sample of strontium is made up of only three isotopes: 86 Sr, 87 Sr and 88 Sr This sample contains 83.00% by mass of 88 Sr This sample of strontium has $A_r = 87.73$

Calculate the percentage abundance of each of the other two isotopes in this sample.

[4 marks]

% abundance ⁸⁷Sr = _____ % abundance ⁸⁶Sr = ____



0 2 . 6

Mg(OH)₂ is used as an antacid to treat indigestion.

A student does an experiment to determine the percentage by mass of Mg(OH)2 in an indigestion tablet.

40.0 cm³ of 0.200 mol dm⁻³ HCl (an excess) is added to 0.200 g of a powdered tablet. The mixture is swirled thoroughly.

All of the Mg(OH)₂ reacts with HCl as shown.

$$Mg(OH)_2 + 2HCl \rightarrow MgCl_2 + 2H_2O$$

The amount of HCl remaining after this reaction is determined by titration with 0.100 mol dm⁻³ NaOH

29.25 cm³ of 0.100 mol dm⁻³ NaOH are needed.

Calculate the percentage by mass of Mg(OH)₂ in the indigestion tablet.

[6 marks]

Percentage by mass

Turn over ▶



| 0 3 | A student uses this method to prepare a standard solution of sodium carbonate. |
|-------|--|
| | Weigh a clean, dry, empty container on a balance that reads to 2 decimal places. Add about 2.5 g of solid sodium carbonate to the container. Tip the solid into a beaker. Add approximately 100 cm³ of distilled water to the beaker and stir until all the solid has dissolved. |
| | 5. Pour the solution into a 250 cm³ volumetric flask. 6. Add distilled water until the top of the meniscus is level with the graduation mark. |
| 0 3.1 | Suggest three improvements to this method. [3 marks] |
| | 1 |
| | |
| | 2 |
| | |
| | 3 |
| | |
| 0 3.2 | A different student uses the correct method to prepare 250 cm³ of sodium carbonate solution in a volumetric flask. The uncertainty for the volumetric flask is ±0.20 cm³ |
| | Calculate the percentage uncertainty in the volume of this sodium carbonate solution. [1 mark] |
| | |
| | |
| | Percentage uncertainty |
| | |
| | |
| | |
| | |
| | |
| | |
| | |



M is a Group 2 metal that forms the nitrate $M(NO_3)_2$ 0.320 g of $M(NO_3)_2$ is heated strongly and decomposes completely.

$$2\, \text{M}(NO_3)_2(s) \ \to \ 2\, \text{MO}\,(s) \ + \ 4\, NO_2(g) \ + \ O_2(g)$$

The mixture of gases formed has a volume of 225 cm3 at 450 °C and 101 000 Pa

Determine the M_r of $M(NO_3)_2$

Identify M.

The gas constant, $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$

[5 marks]

 $M_{\rm r}$ of $\mathbf{M}({\rm NO_3})_2$

Identity of **M**

5

Turn over ▶



| 0 5 | This question is about the shapes of molecules. | |
|-----|--|-----------|
| | Discuss the difference between the shapes of CF ₄ and XeF ₄ In your answer you should: • name the shape of each molecule • explain the shape of each molecule • explain the bond angle(s) in each molecule. | |
| | | [6 marks] |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |



Do not write outside the box

| | ' |
|--|----------|
| | |
| | |






| 0 6 | This question is about halogens and halide ions. | |
|-------|---|-----------|
| 0 6.1 | Explain why the electronegativity of the halogens decreases down the group | [2 marks] |
| | | |
| | | |
| | | |
| | | |
| | Concentrated sulfuric acid reacts with solid sodium chloride and with solid sodium bromide. | |
| 0 6.2 | State one similarity in, and one difference between, these reactions. | [2 marks] |
| | Similarity | |
| | | |
| | Difference | |
| | | |
| 0 6.3 | Solid sodium iodide reacts with concentrated sulfuric acid to form hydrogen | sulfide. |
| | Give a half-equation to show the oxidation of iodide ions. | |
| | Give a half-equation to show the reduction of concentrated sulfuric acid to hydrogen sulfide. | |
| | Use your half-equations to deduce an overall equation for this reaction. | [3 marks] |
| | Half-equation 1 | |
| | | |
| | Half-equation 2 | |
| | | |
| | Overall equation | |
| | | |
| | | |



Do not write outside the







This question is about time of flight (TOF) mass spectrometry.

0 7 . 1

Compound ${\bf X}$ is dissolved in a polar, volatile solvent and is ionised by electrospray ionisation.

Each ion is accelerated so that it has a kinetic energy of $1.36 \times 10^{-16} \, \mathrm{J}$

The kinetic energy of an ion is given by the equation $KE = \frac{1}{2}mv^2$ where:

KE = kinetic energy / J m = mass / kg v = speed / m s⁻¹

The time of flight along the 0.750 m flight tube is 2.48×10^{-5} s

Determine the mass, in g, of one mole of X.

The Avogadro constant, $L = 6.022 \times 10^{23} \text{ mol}^{-1}$

[5 marks]

Mass of one mole of **X**



找名校导师,用小草线上辅导(微信小程序同名)

| Do not writ |
|-------------|
| box |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| - |
| |

| | A mixture of gases is analysed using TOF mass spectrometry. The mixture contains argon, carbon dioxide, nitrogen and oxygen. The mixture is ionised by electron impact. |
|---------|---|
| 0 7.2 | State the meaning of the term electron impact ionisation. [1 mark] |
| 0 7.3 | Identify the ion formed from this mixture that reaches the detector last. |
| | Justify your answer. [2 marks] Ion that reaches detector last |
| | Justification |
| 0 7 . 4 | State how the ions are detected, and how the abundance of each ion is measured, in |
| | a TOF mass spectrometer. [2 marks] How ions are detected |
| | How abundance is measured |
| | |

Turn over for the next question



Find Personal Tutor from www.wisesprout.co.uk 找名校导师,用小草线上辅导(微信小程序同名)

0 8

This question is about the equilibrium mixture formed when **A** and **B** react.

$$A(aq) + 2B(aq) \rightleftharpoons C(aq)$$
 $\Delta H = -32 \text{ kJ mol}^{-1}$

0 8 . 1

A solution containing 0.60 mol of $\bf A$ is added to a solution containing 0.60 mol of $\bf B$. The amount of $\bf C$ formed at equilibrium is 0.28 mol

Deduce the amounts, in moles, of **A** and **B** in this mixture at equilibrium.

[2 marks]

| Amount of A | mo |
|--------------------|----|

Amount of **B** mol

 $\boxed{\mathbf{0} \ \mathbf{8}}$. $\boxed{\mathbf{2}}$ Give an expression for the equilibrium constant (K_c) for this reaction.

[1 mark]

 K_{c}



| 0 8 . 3 | The temperature of the equilibrium mixture is decreased. | | |
|---------|--|--|--|
| | Predict the effect, if any, on the value of $K_{\rm c}$ Give a reason for your prediction. [3 marks] | | |
| | Prediction | | |
| | Reason | | |
| | | | |
| | | | |
| | | | |
| 0 8 . 4 | In another mixture at equilibrium | | |

 $[A] = 0.48 \text{ mol dm}^{-3}$ $[C] = 0.62 \text{ mol dm}^{-3}$

For this reaction, the equilibrium constant K_c = 7.8 mol⁻² dm⁶

Calculate [B] at equilibrium.

Give your answer to the appropriate number of significant figures.

[3 marks]

| B] | mol dm- |
|------------|---------|
| | |



Do not write outside the

Section B

Answer all questions in this section.

Only **one** answer per question is allowed.

For each question completely fill in the circle alongside the appropriate answer.

CORRECT METHOD



WRONG METHODS



If you want to change your answer you must cross out your original answer as shown.



If you wish to return to an answer previously crossed out, ring the answer you now wish to select as shown.

You may do your working in the blank space around each question but this will not be marked. Do not use additional sheets for this working.

0 9 Which atom contains the most neutrons?

[1 mark]

- A 54Cr
- B 55Mn
- **C** 57Fe
- **D** 58Ni





Use this information for Questions 10 and 11.

A student completes a titration to determine the concentration of ethanoic acid in vinegar.

25.0 cm³ of vinegar are transferred to a conical flask using a pipette.

A few drops of phenolphthalein are added to the conical flask.

Sodium hydroxide solution is added from a burette to the conical flask.

The titration is repeated until concordant results are obtained.

1 0 Which suggestion improves the accuracy of the titres?

[1 mark]

- A Rinsing the conical flask with vinegar between each titration.
- ach 👝
- Rinsing the conical flask with sodium hydroxide solution between each titration.



0

C Rinsing the conical flask with water between each titration.

| Γ | 0 | |
|---|---|--|

D Not rinsing the conical flask between each titration.

1 1 Which suggestion decreases the percentage uncertainty in the mean titre?



A Use a more dilute solution of sodium hydroxide in the burette.



B Use a more dilute solution of vinegar.



c Rinse the inside of the conical flask with distilled water during each titration.

| 0 |
|---|

Rinse the tip of the burette with distilled water near the end point in

| | $\overline{}$ | |
|---|---------------|---|
| - | | _ |
| | | |

each titration.

| 0 | |
|---|--|
| | |

1 2 Which reaction has the highest percentage atom economy for the production of hydrogen?

[1 mark]

A LiH + $H_2O \rightarrow LiOH + H_2$



 $\textbf{B} \ CO \ + \ H_2O \ \rightarrow \ CO_2 \ + \ H_2$



- **C** 2Al + $3H_2O \rightarrow Al_2O_3 + 3H_2$
- **D** $CH_4 + H_2O \rightarrow CO + 3H_2$
- 0

Turn over for the next question





A NCl₃

Which molecule has a permanent dipole?

B CCl₄

C PF₅

D SF₆

Which molecule can accept an electron pair during the formation of a coordinate bond?

[1 mark]

[1 mark]

A NH₃

B AlCl₃

C SiH₄

D PCl₃

1 5 Which reaction has an enthalpy change equal to the standard enthalpy of formation of potassium oxide?

[1 mark]

 $A \ \, 4\,K(s) \,\, + \,\, O_2(g) \,\, \to \,\, 2\,K_2O(s)$

0

 $\textbf{B} \ 2 \, \text{K(s)} \ + \ \text{O(g)} \ \rightarrow \ \text{K}_2 \text{O(s)}$

0

C $2K^+(g) + O^{2-}(g) \rightarrow K_2O(s)$

0

 $\mbox{\bf D} \ \, 2 \, K(s) \, \, + \, \, \frac{1}{2} \, O_2(g) \, \, \to \, \, K_2O(s)$



1 6 In w

In which species is chlorine in its highest oxidation state?

[1 mark]

- A ClF₂-
- B ClO₄-
- C ClO₂
- D ClF₃

1 7 Which statement about this redox reaction is correct?

$$3\,Sn^{2+}(aq) \ + \ Cr_2O_7{}^{2-}(aq) \ + \ 2\,H^+(aq) \ \rightarrow \ 2\,Cr^{3+}(aq) \ + \ 3\,SnO_2(s) \ + \ H_2O(I)$$

[1 mark]

- **A** Sn²⁺ is the oxidising agent and it gains electrons.

 □
- **B** Sn²⁺ is the reducing agent and it gains electrons.
- **C** $Cr_2O_7^{2-}$ is the oxidising agent and it gains electrons.
- **D** $Cr_2O_7^{2-}$ is the reducing agent and it gains electrons.

Which incomplete half-equation is balanced by adding two H⁺ ions and one electron to the left-hand side?

[1 mark]

- **A** $CH_3CHO \rightarrow CH_3CH_2OH$
- **B** $VO^{2+} \rightarrow V^{3+} + H_2O$
- **C** $HNO_2 \rightarrow NO + H_2O$
- $\mathbf{D} \ \mathsf{O}_2 \to \mathsf{H}_2\mathsf{O}_2$

Turn over for the next question



Use this information for Questions 19 to 23.

A student completes some test-tube reactions on five solutions, **P**, **Q**, **R**, **S** and **T**. The student completes each test on separate samples of each solution. Observations are shown in the table.

| Solution | Test 1 Add a few drops of H₂SO₄(aq) | Test 2 Add HNO ₃ (aq) then a few drops of AgNO ₃ (aq) | Test 3 Add a few drops of NaOH(aq) | Test 4 Add a few drops of Cl ₂ (aq) |
|----------|-------------------------------------|---|--|--|
| Р | White precipitate | Cream precipitate | No visible change | Orange solution |
| Q | Effervescence | Effervescence and white precipitate | No visible change | No visible change |
| R | No visible change | No visible change | White precipitate | No visible change |
| s | White precipitate | | No visible change | Dark brown solution |
| Т | No visible change | White precipitate | White precipitate | No visible change |

Use the information in the table to answer Questions 19 to 23.


| 1 9 | Which solution co | ontains carbonate ions? | [1 mark] |
|-----|----------------------------|---|----------|
| | A Solution P | 0 | |
| | B Solution Q | 0 | |
| | C Solution R | 0 | |
| | D Solution S | 0 | |
| 2 0 | What could be the | e identity of the compound in solution P ? | [1 mark] |
| | A MgBr ₂ | 0 | |
| | B BaBr ₂ | 0 | |
| | C MgCl ₂ | 0 | |
| | D BaCl ₂ | 0 | |
| | | | |



| | 名校导师,用小草线上辅导 |
|----|---------------|
| |) 小難二 |
| | (微信小桂序同名 |
| 15 | 予回名 |
| | $\overline{}$ |

| 2 1 | What observation is expected in Test 2 for solution S ? | [1 mark] | |
|-----|---|----------|--|
| | A No visible change | | |
| | B Effervescence | | |
| | C White precipitate | | |
| | D Pale yellow precipitate | | |
| 2 2 | In Test 2 , the mixture formed by solution S is filtered. | | |
| | Which substance is present in the filtrate? | [1 mark] | |
| | A AgBr | | |
| | B Agl | | |
| | C Ba(NO ₃) ₂ | | |
| | D Mg(NO ₃) ₂ | | |
| 2 3 | What could be the identity of the compound in solution T ? | [1 mark] | |
| | A MgCl ₂ | | |
| | B BaCl ₂ | | |
| | C MgBr ₂ | | |
| | D BaBr ₂ | | |
| | END OF QUESTIONS | | |







Do not write outside the box

| •••••• |
|--|
| |
| ••••• |
| ••••• |
| •••••• |
| ·••••••••••••••••••••••••••••••••••••• |
| •••••• |
| •••••• |
| •••••• |
| •••••• |
| •••••• |
| •••••• |
| ••••••• |
| •••••• |
| ······································ |
| · ······ |
| ••••••• |
| •••••• |
| ······································ |
| |
| |
| |
| |
| · ····· |



| Question number | Additional page, if required. Write the question numbers in the left-hand margin. |
|-----------------|--|
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | Copyright information For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet |
| | is published after each live examination series and is available for free download from www.aqa.org.uk Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the |
| | Copyright Team. Copyright © 2023 AQA and its licensors. All rights reserved. |



