Please check the examination deta	ils belo	w before ente	ring your candidate information
Candidate surname			Other names
Pearson Edexcel Level 1/Level 2 GCSE (9–1)	Cent	re Number	Candidate Number
Time 1 hour 10 minutes		Paper reference	1SC0/2CH
Combined Scient PAPER 5 Higher Tier	nce	2	
You must have: Calculator, ruler			Total Marks

### **Instructions**

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions.
- Answer the questions in the spaces provided
  - there may be more space than you need.
- Calculators may be used.
- Any diagrams may NOT be accurately drawn, unless otherwise indicated.
- You must show all your working out with your answer clearly identified at the end of your solution.

### Information

- The total mark for this paper is 60.
- The marks for **each** question are shown in brackets
  - use this as a guide as to how much time to spend on each question.
- In questions marked with an **asterisk** (\*), marks will be awarded for your ability to structure your answer logically showing how the points that you make are related or follow on from each other where appropriate.
- There is a periodic table on the back cover of the paper.

### **Advice**

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- Good luck with your examination.

Turn over ▶







# Answer ALL questions. Write your answers in the spaces provided.

Some questions must be answered with a cross in a box  $\boxtimes$ . If you change your mind about an answer, put a line through the box  $\boxtimes$  and then mark your new answer with a cross  $\boxtimes$ .

1 (a) The concentration of a solution can be calculated using the equation

concentration of solution = 
$$\frac{\text{mass of solid}}{\text{volume of solution}}$$

A student dissolved 9.25 g of ammonium chloride in water and made up the solution to a volume of 200 cm<sup>3</sup>.

Use the equation to calculate the concentration of this solution in  $g\,dm^{-3}$ .

(2)

.....

(b) Dissolving ammonium chloride in water is an endothermic process. Figure 1 shows part of the reaction profile for this process.

heat energy

ammonium chloride solution

ammonium chloride + water

progress of reaction

## Figure 1

(i) Explain how Figure 1 shows that dissolving ammonium chloride in water is an endothermic process.


(2)



(ii) Complete the reaction profile in Figure 1 and label the activation energy.

(2)

(c) A student used the equipment in Figure 2 to investigate whether electricity can pass through solid ammonium chloride and through ammonium chloride solution.






Figure 2

If an electrical current flows in the circuit, the lamp will light up.

Figure 3 shows the results of the investigation.

substance	lamp
solid ammonium chloride	did not light up
ammonium chloride solution	lit up brightly

Figure 3

<b>Explain</b>	the	results	of the	investic	iation.
----------------	-----	---------	--------	----------	---------

(3)

(Total for Question 1 = 9 marks)

2	Diesel oil is a	a mixture of h	ydrocarbons	that can	be obtained	from crude oil.	

(a) State the name of the process used to separate diesel oil from crude oil.

(1)

(b) Diesel oil contains alkanes.

These alkanes are part of an homologous series.

Which statement about compounds in this homologous series is true?

(1)

- A they have the same chemical formula
- **B** they have the same empirical formula
- C they have the same general formula
- D they have the same molecular formula
- (c) When fuels such as diesel oil are burned, the high temperatures produced can cause nitrogen and oxygen in the air to form the pollutant nitrogen dioxide.

Complete the balanced equation for the reaction.

(2)

$$N_2 + 2O_2 \rightarrow \dots$$

(d) Explain how the greenhouse effect is caused by the gases produced by the complete combustion of diesel oil.

(3)

(Total for Question 2 = 7 marks)



3	This gues	tion	is about notacsium and zinc	
3			is about potassium and zinc.	_
	(a) Which	n of t	he following temperatures is most likely to be the melting po	oint of potassium? (1)
	$\times$	A	-63°C	
	X	В	6.3°C	
	X	C	63°C	
	X	D	630°C	
			w the electronic configuration of an atom of potassium is related to the periodic table.	ated to its (2)
	(c) Potas	sium	reacts with oxygen to form potassium oxide.	
	(i) D	escri	oe the test to show that a gas is oxygen.	(2)
	(ii) Po	otass	ium oxide is ionic.	
	W	rite <sup>·</sup>	he electronic configurations for the ions in potassium oxide,	K <sub>2</sub> O. (2)
			potassium ion:	
			oxide ion:	

(d) Figure 4 shows two gas syringes connected by a glass tube.

Inside the glass tube there are some pieces of zinc. Zinc reacts with oxygen at a temperature of over 225 °C. Not all the oxygen reacts at once, the oxygen reacts only when in contact with the zinc.





Figure 4

Devise a plan to find the volume of oxygen contained in a known volume of air, using the apparatus shown in Figure 4.



**4** This question is about the rate of reaction between calcium carbonate and dilute hydrochloric acid.

The word equation for this reaction is

calcium carbonate + hydrochloric acid → calcium chloride + water + carbon dioxide

(a) Which of the following is the formula for calcium carbonate?

(1)

- A CaCO₂
- B CaCO<sub>3</sub>
- C Ca(CO)<sub>3</sub>
- **D** Ca(CO<sub>2</sub>)<sub>2</sub>
- (b) Some pieces of calcium carbonate were added to dilute hydrochloric acid in a conical flask and the volume of carbon dioxide produced was measured.

Complete the diagram in Figure 5 to show the apparatus to collect the gas produced and measure its volume.

(2)




Figure 5

- (c) The reaction between calcium carbonate and dilute hydrochloric acid was investigated at different temperatures.
  - (i) State what could be used to keep the temperature of the conical flask and its contents at a temperature of 45 °C throughout the reaction.

(1)



(ii) Figure 6 shows a graph of volume of gas collected in this investigation.

volume of gas in cm<sup>3</sup>




Figure 6

Draw a tangent at 100 seconds on Figure 6. Use this tangent to calculate the rate of reaction at this time.

(2)

rate of reaction = .... cm<sup>3</sup> s<sup>-1</sup>

(iii) The temperature of the acid was kept at 45 °C.  State <b>one</b> other variable that needs to be controlled during this investigation.	(1)
(iv) Explain, in terms of particles, how decreasing the temperature affects the rate of this reaction.	(3)
(Total for Question 4 = 10 ma	rks)

10

- 5 This question is about some of the elements in group 7 of the periodic table.
  - (a) Which row in the table correctly shows the colours and physical states of the elements at room temperature?

(1)

×	Α	iodine: purple gas	bromine: yellow liquid
X	В	chlorine: pale green gas	iodine: brown solid
×	C	bromine: red-brown liquid	chlorine: yellow liquid
×	D	iodine: dark grey solid	bromine: red-brown liquid

(b) The compound phosphorus oxychloride has the formula POCl<sub>3</sub>.

Calculate the percentage by mass of chlorine in phosphorus oxychloride.

(relative atomic masses: O = 16.0, P = 31.0, Cl = 35.5)

(2)

percentage by mass of chlorine = .....

(c) When iron reacts with chlorine, iron chloride is formed.

Two possible equations for this reaction are

**A** Fe + 
$$Cl_2 \rightarrow FeCl_2$$

**B** 
$$2Fe + 3Cl_2 \rightarrow 2FeCl_3$$

In an experiment, 8.40 g iron reacts with chlorine to form 19.05 g iron chloride.

Show, using a calculation, which reaction, **A** or **B**, is taking place.

You must show your working.

(relative atomic masses: Cl = 35.5, Fe = 56.0)

(3)



\*(d) Group 1 metals react with the elements from group 7 to form salts.

Some examples of these reactions are shown in Figure 7.

reaction	word equation	
W	lithium + chlorine $ ightarrow$ lithium chloride	
X	potassium + fluorine $ ightarrow$ potassium fluoride	
Y rubidium + iodine → rubidium iodide		
z	potassium + bromine → potassium bromide	

## Figure 7

You will find the position of these elements in their groups on the periodic table.

Explain, in terms of their electronic configurations and the relative reactivity of these elements, which of the reactions shown in Figure 7 would be the most violent.

(6)

(Tota	for Question 5 = 12 marks)



- **6** Pentadecane, C<sub>15</sub>H<sub>32</sub>, is a hydrocarbon and is used as a fuel.
  - (a) The incomplete combustion of pentadecane produces carbon monoxide. Carbon monoxide is a toxic gas.
    - (i) Explain why the incomplete combustion of pentadecane can produce carbon monoxide as one of the products.

(2)

(ii) Explain how carbon monoxide behaves as a toxic gas.

(2)

(b) 1 mole of pentadecane can be cracked to form 1 mole of octane, C<sub>8</sub>H<sub>18</sub>, and 1 mole of propene, C<sub>3</sub>H<sub>6</sub>, and 2 moles of another product.

Complete the balanced equation for this reaction by adding the formula of the missing product.

(1)

$$C_{15}H_{32} \rightarrow C_8H_{18} + C_3H_6 + 2$$



(c) Figure 8 shows the reaction of propene,  $C_3H_6$ , with water.

Figure 8

Figure 9 shows some bond energies.

bond	bond energy in kJ mol <sup>-1</sup>
С—С	347
C—O	358
С—Н	413
О—Н	464
C=C	612

Figure 9

Use the bond energies in Figure 9 to calculate the energy change of the reaction in Figure 8.

•••••	 	 	 	 	 	 	

energy change of reaction = .....kJ mol<sup>-1</sup>



(4)

(2)

(d) Methane gas,  $CH_4$ , was burned using the apparatus shown in Figure 10.




Figure 10


(Total for Question 6 = 11 marks)

**TOTAL FOR PAPER = 60 MARKS** 





# The periodic table of the elements

0	4 <b>He</b> helium 2	20 <b>Ne</b> neon 10	40 <b>Ar</b> argon 18	84 <b>Kr</b> krypton 36	131 <b>Xe</b> xeron 54	[222] <b>Rn</b> radon 86
7		19 <b>F</b> fluorine 9	35.5 <b>CI</b> chlorine 17	80 <b>Br</b> bromine 35	127 	[210] <b>At</b> astatine 85
9		16 <b>0</b> oxygen 8	32 <b>S</b> sulfur 16	79 <b>Se</b> selenium 34	128 <b>Te</b> tellurium 52	[209] <b>Po</b> polonium 84
2		14 <b>N</b> nitrogen 7	31 P phosphorus 15	75 <b>As</b> arsenic 33	122 <b>Sb</b> antimony 51	209 <b>Bi</b> bismuth 83
4		12 <b>C</b> carbon 6	28 <b>Si</b> silicon 14	73 <b>Ge</b> germanium 32	119 <b>Sn</b> tin 50	207 <b>Pb</b> lead 82
က		11 <b>B</b> boron 5	27 Aluminium 13	70 <b>Ga</b> gallium 31	115 In indium 49	204 <b>T</b> thallium 81
	·			65 <b>Zn</b> zinc 30	112 <b>Cd</b> cadmium 48	201 <b>Hg</b> mercury 80
				63.5 <b>Cu</b> copper 29	108 <b>Ag</b> silver 47	197 <b>Au</b> gold 79
				59 <b>Ni</b> nickel 28	106 <b>Pd</b> palladium 46	195 <b>Pt</b> platinum 78
				59 <b>Co</b> cobalt 27	103 <b>Rh</b> rhodium 45	192 <b>Ir</b> iridium 77
	1 <b>H</b> hydrogen				101 <b>Ru</b> ruthenium 44	190 <b>Os</b> osmium 76
•				55 Mn manganese 25	[98] <b>Tc</b> technetium 43	186 <b>Re</b> rhenium 75
		mass <b>bol</b> lumber		52 Cr chromium 24	96 <b>Mo</b> molybdenum 42	184 <b>W</b> tungsten 74
	Key	relative atomic mass atomic symbol name atomic (proton) number		51 V vanadium 23	93 <b>Nb</b> niobium 41	181 <b>Ta</b> tantalum 73
		relati <b>atc</b> atomic		48 <b>Ti</b> titanium 22	91 <b>Zr</b> zirconium 40	178 <b>Hf</b> hafnium 72
				45 Sc scandium 21	89 <b>Y</b> yttrium 39	139 <b>La</b> * lanthanum 57
2		9 <b>Be</b> beryllium 4	24 <b>Mg</b> magnesium 12	40 <b>Ca</b> calcium 20	88 <b>Sr</b> strontium 38	137 <b>Ba</b> barium 56
_		7 <b>Li</b> lithium 3	23 <b>Na</b> sodium 11	39 <b>K</b> potassium 19	85 <b>Rb</b> rubidium 37	133 <b>Cs</b> caesium 55

<sup>\*</sup> The elements with atomic numbers from 58 to 71 are omitted from this part of the periodic table.

The relative atomic masses of copper and chlorine have not been rounded to the nearest whole number.